

2002

Subjective Assessment of Sleep Quality Across the Menstrual Cycle in Women with Premenstrual Dysphoric Disorder

Jonathan Bruce Santo

University of Nebraska at Omaha, jsanto@unomaha.edu

E. Chevrier

McGill University

P. L'Esperance

McGill University

D. B. Boivin

McGill University

Follow this and additional works at: <https://digitalcommons.unomaha.edu/psychfacpub>

 Part of the [Psychology Commons](#)

Recommended Citation

Santo, Jonathan Bruce; Chevrier, E.; L'Esperance, P.; and Boivin, D. B., "Subjective Assessment of Sleep Quality Across the Menstrual Cycle in Women with Premenstrual Dysphoric Disorder" (2002). *Psychology Faculty Publications*. 15.

<https://digitalcommons.unomaha.edu/psychfacpub/15>

This Article is brought to you for free and open access by the Department of Psychology at DigitalCommons@UNO. It has been accepted for inclusion in Psychology Faculty Publications by an authorized administrator of DigitalCommons@UNO. For more information, please contact unodigitalcommons@unomaha.edu.



717.Q

SUBJECTIVE ASSESSMENT OF SLEEP QUALITY ACROSS THE MENSTRUAL CYCLE IN WOMEN WITH PREMENSTRUAL DYSPHORIC DISORDER

Santo JB,¹ Chevrier E,¹ L'Esperance P,¹ Boivin DB¹

⁽¹⁾ Douglas Hospital Research Center, Department of Psychiatry, McGill University

Introduction: Existing evidence demonstrates that sleep structure varies across the menstrual cycle in healthy women (1). These variations could be more severe in women suffering from premenstrual dysphoric disorder (PMDD) (2). In a previous study of healthy women, subjective sleep quality was shown to be constant across both phases of the menstrual cycle (3). The current study aims to test whether there exists a variation in subjective sleep quality of PMDD sufferers across the follicular and luteal phases of the menstrual cycle.

Methods: Five women (ages 28-41) with a clinical diagnosis of PMDD maintained regular sleep/wake habits over one entire menstrual cycle. Sleep/wake times outside of the laboratory were kept constant and confirmed by wrist actigraphy. Daytime napping was prohibited during the entire study. Every third night of one entire menstrual cycle, sleep was polysomnographically recorded in the laboratory using a standard montage. Intrinsic sleep disorders were ruled out during the first sleep recording. Participants also assessed the quality of their prior sleep episode, upon waking using a 10 cm visual analog scale.

Results: Mean sleep length in the laboratory was 8h12min + 10 minutes. A Wilcoxon signed ranks test was performed to compare subjective sleep quality scores of the early follicular (first week) and the late luteal (last week) phases of the menstrual cycle. During the early follicular phase, subjective sleep quality was 6.43 (SD= 2.89), while that of the late luteal phase was 5.41 (SD = 3.43) (Figure 1). The scores were found to vary significantly between the phases of the menstrual cycle ($p = 0.041$). Furthermore, participants' assessment of sleep quality was positively correlated with polysomnographic measures of sleep efficiency ($r = 0.550$, $p = 0.02$) (Figure 2).

Figure 1





Figure 2



Conclusions: The present study indicates that subjective sleep quality deteriorates significantly during the late luteal phase in women with PMDD. This is not surprising since sleep disturbances are an important diagnostic criteria of PMDD. These results are different from what has been reported in healthy women (3). Elucidation of the factors mediating changes in sleep during menstruation in participants with PMDD may have key implications in developing new and innovative treatments.

References:

- (1) Driver HS, & Baker FC. Menstrual Factors in Sleep. *Sleep Medicine Reviews* 1998;2:213-229.
- (2) Mauri M, Reid RL & MacLean AW. Sleep in the Premenstrual Phase: a Self Report Study of PMS Patients and Normal Controls. *Acta Psychiatrica Scandinavia* 1988;78:82-86.
- (3) Driver HS, Dijk DJ, Werth E, et al. Sleep and the Sleep Electroencephalogram across the Menstrual Cycle in Young Healthy Women. *Journal of Clinical Endocrinology and Metabolism* 1996;81:728-735.

Research supported by an operating grant from the Canadian Institutes of Health Research.