

University of Nebraska at Omaha DigitalCommons@UNO

Health and Kinesiology Faculty Publications

School of Health and Kinesiology

9-2017

Diagnosis of a Chiari Malformation After a Concussion in a Junior College Football Player With a History of Chronic Headaches: A Case Report

Takeaki Ando University of Nebraska at Omaha

Shannon Gehr Iowa Western Community College

Melanie L. McGrath The University Of Montana

Adam B. Rosen University of Nebraska at Omaha, arosen@unomaha.edu

Follow this and additional works at: https://digitalcommons.unomaha.edu/hperfacpub

Part of the Health and Physical Education Commons, and the Kinesiology Commons Please take our feedback survey at: https://unomaha.az1.qualtrics.com/jfe/form/ SV_8cchtFmpDyGfBLE

Recommended Citation

Ando, Takeaki; Gehr, Shannon; McGrath, Melanie L.; and Rosen, Adam B., "Diagnosis of a Chiari Malformation After a Concussion in a Junior College Football Player With a History of Chronic Headaches: A Case Report" (2017). *Health and Kinesiology Faculty Publications*. 30. https://digitalcommons.unomaha.edu/hperfacpub/30

This Article is brought to you for free and open access by the School of Health and Kinesiology at DigitalCommons@UNO. It has been accepted for inclusion in Health and Kinesiology Faculty Publications by an authorized administrator of DigitalCommons@UNO. For more information, please contact unodigitalcommons@unomaha.edu.



1	Diagnosis of a Chiari Malformation after a Concussion in a Junior College Football Player with a
2	history of Chronic Headaches: A Case Report
3	
4	Key Points:
5	
6	- Junior college football player who was diagnosed post-concussion with a Chiari
7	malformation.
8	- Chiari malformations are frequently missed on images which can complicate treatments.
9	- Treatment protocols for those with a concussion with a Chiari malformation are debatable.
10	
11	
12	Key Words: brain, return to play, head injury, sport-related, collision sport
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	

23 ABSTRACT

24	The purpose of this report is to present the case of a National Junior Collegiate Athletic
25	Association football player diagnosed with Chiari malformation post-concussion. A Chiari
26	malformation is characterized by the cerebellum presenting below the level of the foramen. The
27	uniqueness of this case stems from the patients' health history, length of symptoms and
28	diagnosis. The effectiveness of treatment options, and the primary means to reduce the risk of
29	catastrophic head injury in those with Chiari Malformations are debatable. Clinicians should be
30	familiar with the potential for the presence of a Chiari malformation with persistent symptoms
31	post-concussion.
32	
33	
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	

45 INTRODUCTION

46	Concussions are a significant issue in athletics with an estimated 1.6-3.8 million
47	individuals suffering from concussion while performing physical activities each year in the
48	United States. ^{1,2} Abnormal brain anatomical structures, while uncommon, may play a significant
49	role in concussion management. ³⁻⁵
50	Chiari malformation, also known as Arnold-Chiari malformation, is a rare anatomical
51	irregularity in which the cerebellum may be depressed 3-5mm below the level of the foramen
52	magnum. ⁶ Onset of Chiari malformation may be congenital or acquired through infection or
53	trauma and is only diagnosed between 0.1-3.6% of all people, possibly because of its
54	asymptomatic nature. ⁶⁻¹¹ Chiari malformations are normally diagnosed through magnetic
55	resonance imaging (MRI), and a radiological measurement of the amount of cerebellar
56	depression. ^{6,7} There are four types of this deformity, however, a Type I is the most common
57	variant and involves downward displacement of the cerebellar tonsils emanating from the
58	inferior opening of the skull and into the spinal cord without displacing the other structures of the
59	central nervous system.
60	The symptoms of Chiari malformations are similar to post-concussion syndrome, and
61	include headache, dizziness, muscle weakness, chronic fatigue, tinnitus and impaired ability to
62	coordinate movement. Those symptoms can be aggravated by a Valsalva maneuver, cough, or
63	postural changes, and may derive from disturbing cerebrospinal fluid or cerebellum compression
64	through trauma. ^{6,8,9,11} However, no studies have clearly demonstrated a specific age of onset of
65	Chiari malformation in acquired cases. ^{6,8,9,11} It is generally believed that surgery is a superior
66	treatment option compared to conservative treatments, but is dependent on the physicians'

67 impression of the severity and associated symptoms.^{6,7,8,12,13}

Due to its' asymptomatic presentation, athletes may have a Chiari malformation, but not
be diagnosed until an MRI is performed post-concussion with severe or unresolving
symptoms.^{6,8-11,14,15} Therefore, the purpose of this case report is to present the unique case of a
National Junior Collegiate Athletic Association football player who was diagnosed with a Type I
Chiari malformation after a concussion.

73 CASE PRESENTATION

74 Patient

The patient was a male Junior college student and football linebacker (Caucasian, 18 75years, 182cm, 104.3kg). He had no outstanding previous injuries related, but reported headaches 76almost daily for approximately three years and noticed a steady increase in headache frequency. 77The headaches were mainly posterior and radiated forward to the superior portion his head and 78forehead, and he also reported pressure behind his eyes during these episodes. He tolerated most 7980 of the headaches, but indicated several were severe. A severe headache for the patient was characterized as global with pressure behind his eyes, photophobia, phonophobia and occasional 81 nausea. There were no other neurologic features or consistent length of symptoms. At the age of 82 83 16, his headaches worsened and an MRI of the brain was read as normal by a neurologist.

During a football practice on Aug 13th 2012, he tackled and hit heads with an opposing player. He was evaluated immediately by the certified athletic trainer (AT). The athlete did not remember how he got hit and could not answer questions about the incident correctly. The player was asked his name, but he answered with a different teammate's name. His symptoms included headache, vomiting, dizziness, visual problems, fatigue, photophobia, sensitivity to noise,

4

89	retrograde amnesia, emotional problems, forgetfulness, positive Romberg test, and difficulty
90	walking. The athlete was referred to the local emergency room where he was diagnosed with a
91	grade II concussion. Moreover, cranial CT scans revealed no acute findings but showed a
92	possible instance of Type I Chiari malformation.
93	During the week following the injury, the AT checked on his status and symptoms daily.
94	His symptoms slowly improved, however he performed an ImPACT (ImPACT Applications, Inc.
95	Version 2.1, Pittsburgh, PA, USA) test which showed each score worse than his baseline (Table
96	1). Approximately one week later, he was seen by a ^{1st} neurologist. The 1 st neurologist diagnosed
97	him with post-concussive syndrome due to his persistent headaches and occasional nausea. Upon
98	evaluation by the 1 st neurologist he demonstrated errors in serial-7 subtractions and occasional
99	delays in information retrieval. Language and speech, eye movement, facial sensation, muscle
100	strength assessments, and coordination tests were normal. An ImPACT test one week later, while
101	improved still demonstrated worse than baseline scores.
102	He was seen again by the 1 st neurologist approximately three weeks later. The patient
103	displayed improvements in cognitive memory function, but noticed the headaches had not
104	ceased. In addition, he reported the headaches were so severe, he had very little appetite. The 2 nd
105	neurologist reviewed his previous MRI as well as a CT scan of his head, and diagnosed him with
106	Type I Chiari malformation with chronic headaches. The 2 nd neurologist found a 3-mm
107	herniation on his MRI (Figure 1a). Although the 2 nd neurologist diagnosed him with a Chiari
108	malformation, they believed that the headaches were not related to this disorder because this was
109	a congenital malformation that had not altered over time. Since the patient's headaches had not
110	improved with medication, they recommended that he see a 3 rd neurologist for a second opinion

- 111 on his headaches. He saw the 3rd neurologist approximately a week later, but he could not
- 112 determine the exact factors which led to his symptoms.
- 113 Intervention
- 114 The primary treatment was rest to lessen the symptoms and neurological issues. The 1st
- 115 neurologist recommended that prior to return to play the patient would need to be asymptomatic
- both at rest and during activity. In addition, the ^{1st} and ^{2nd} neurologist suggested no clearance for
- 117 physical activity until his ImPACT score was closer to his baseline. He was also prescribed
- 118 Midrin (Isometheptene, dichloralphenazone, and acetaminophen, 1-2 capsules, TID-QID, prn) as
- 119 a trial therapy for his headaches, which is a typical pharmacological agent for severe migraines.
- 120 After three-weeks from the visit to the 1st neurologist, he was still not cleared to return
- 121 to playing football and continued to report his progress to the AT. The Midrin was discontinued
- 122 after the fourth visit because it appeared to provide no benefit and Nortriptyline (25mg-50mg at
- 123 bedtime), a tricyclic antidepressant and typically used for severe chronic migraines was
- 124 prescribed. The neurologist recommended continuing with conservative care and suggested the
- 125 patient seek a second opinion.
- 126 He sought the second opinion and was seen by a 3rd neurologist just over one month
- 127 from the initial injury due to no significant improvements in headaches or symptoms. The 3rd
- 128 neurologist believed there was a low likelihood that the headaches were related to the Chiari
- 129 malformation. The neurologist started him on Topamax as a preventative measure, and he was
- 130 instructed to take nortriptyline for sleep as needed. He also continued the previously prescribed
- 131 medications. After this consultation, he saw a different neurosurgeon one month later. This 3rd
- 132 neurologist also recommended no sports or physical activities until his symptoms and headaches

133 were resolved.

134The student decided to end his participation on the football team in three months after the initial injury because his symptoms, especially the headaches, had not improved, and the 135136neurologists' opinion had not changed. He was recommended to a 4th neurosurgeon. This neurosurgeon believed his Chiari 137138malformation was symptomatic with 14-mm herniation from the previous MRI. This 4th 139neurosurgeon felt he was a candidate for surgery because of a loss of the gag reflex, a sign of a cranial nerve X problem. He had a surgical intervention 6 months later, with no improvement in 140the preceding months. The 4th neurosurgeon performed a posterior fossa decompression with 141 142external durotomy. Post-surgery, he was prescribed Bacolfen to prevent muscle spasms and Dilaudid for pain control. 143**Comparative Outcome** 144Symptoms began improving in the three months post-surgery (Figure 2). He saw the 145146neurosurgeon three times at six month intervals. His symptoms, including the severe headaches began returning, and at the final follow-up appointment of 18 months post-surgery the 147neurosurgeon told him there was nothing else which could be done. The case remains 148149unresolved. 150Discussion The purpose of this case study was to present a unique report of an athlete that had a 151

152 concussion, which resulted in a Type I Chiari Malformation diagnosis. The uniqueness of this
153 case extends from the health history and length of symptoms of the patient. Moreover, several
154 MRI's were performed on the patient in which initial imaging was read as normal.

155	This case has several implications on clinical practice. First, Chiari malformations are
156	rare and typically diagnosed in infancy, so it is possible that previously undiagnosed conditions
157	in older patients may be overlooked by physicians. Chiari malformations are typically diagnosed
158	by an MRI by measuring the depression of the cerebellum. ⁶⁻¹¹ The initial missed diagnosis likely
159	resulted from the rarity of Chiari malformations and the expertise required to identify them. ^{6,8,9}
160	Therefore, patients at the high school and collegiate levels may be diagnosed during post-
161	concussion imaging investigations. ^{13,14,15} Clinicians should also take this into consideration,
162	when the patient's symptoms do not improve for lengthy periods. ^{6,10,16}
163	There are several symptoms which can overlap in patients with concussion and/or Chiari
164	malformation. Typical symptoms which could be due to both conditions include severe
165	headache, dizziness, fatigue, tinnitus, nausea and an impaired ability to coordinate movement.
166	^{6,8,9,11} Differential symptoms for patients with solely a Chiari malformation may include vagus
167	nerve dysfunction (i.e. difficulty swallowing) as well as numbness and abnormal sensations in
168	the extremities due to the compression on the cerebellum. ^{6,8,9,11} However without the imaging in
169	this particular case, it was exceedingly difficult to differentiate among the diagnoses of post-
170	concussion symptoms, chronic headaches, and Chiari malformation due to the similarity in
171	clinical presentation.
172	Conservative treatments are typically recommended for Chiari Malformations prior to
173	surgical interventions. For this particular case, physical rest and various medications were
174	ineffective for treating the headaches and symptoms. Therefore, this case ultimately turned to a
175	surgical intervention, which some studies have demonstrated better outcomes compared to
176	conservative treatments. ^{7,8,12,13,16,17} Surgical interventions appear to decrease the risk of

8

reoccurrence and improve symptoms including headaches.^{7,8,12,13,16,17} However, the severity of
the condition and the patient and neurosurgeon preference influence the course of treatment and
further research is still necessary to identify the ideal protocol.

180 Similar cases have been reported in the literature including a female volleyball player⁴ and male football players.^{10,14,15} However, each of these athletes were asymptomatic prior to the 181 182concussive episode. In each case, symptoms worsened post-concussion and the neurologists 183 diagnosed them with a Type I Chiari malformation from their MRI. Each did not receive surgery, underwent conservative protocols, recovered, and were able to return to sports without incident. 184In a surgical case, Callaway et al¹⁶ reported a football player suffering from tingling, radiating 185186into both anterior thighs as well as shortness of breath after a contact injury. A follow-up MRI of the patient revealed a Type I Chiari malformation and his post-surgical recovery was uneventful. 187 However, due to the mechanism of injury, several orthopedic and neurosurgical spine specialists 188 advised the patient not to return to contact sports. Based on these published case studies there 189 190appears to be no consensus on the best treatment or return to activity path for athletes with Chiari malformations and depends on the individual patient.^{6-8,10,12-14,16-18} However unlike the previous 191cases, both conservative management and surgical interventions failed, which suggests a need for 192further options for patients. 193194Criteria for returning to competition requires the patient to be asymptomatic and receive physician clearance. It was no different in this particular case and in addition to those 195196 requirements the neurologists wanted the patients ImPACT test scores to be similar to baseline prior to any return to play decision. Albeit the current patient was still experiencing symptoms, 197

198 one study suggests the risk for athletes with asymptomatic Chiari malformation is low while

9

199	playing various kinds of sports including contact sports. ⁹ Clinicians may take this into
200	consideration when deciding whether to allow an athlete to return to play, however a cautious
201	approach is warranted.
202	CLINICAL BOTTOM LINE
203	It is difficult to clinically differentiate between the source of symptoms when both a
204	concussion and Chiari malformation are present as there is a potential for an overlap in
205	symptoms. Clinicians should be familiar with the possibility of a Chiari malformation when
206	patients are suffering from long-term, unrelenting headaches or other symptoms post-concussion.
207	Additionally, with persistent symptoms in individuals diagnosed with a concussion, an MRI may
208	be necessary to rule-out structural abnormalities. Creating imaging guidelines may be helpful for
209	patients following head injuries with persistent or unusual symptoms, so future studies should
210	work to develop specific guidelines for these cases. The effectiveness of treatment options
211	whether surgical or non-surgical, and the best ways to lessen the risk of catastrophic head
212	injuries in those with Chiari Malformations are still largely unknown. Further studies are needed
213	to clarify these issues, and should lead to creating guidelines for athletes with Chiari
214	malformation intended to allow them to participate in sports safely.
215	
216	REFERENCES
217	1. Langlois JA, Rutland-Brown W, Wald MM. The epidemiology and impact of traumatic brain
218	injury: a brief overview. J Head Trauma Rehabil, 2006; 21:375-378.
219	2. Broglio SP, Cantu RC, Gioia GA, et al. National Athletic Trainers' Association Position
220	Statement: Management of Sport Concussion. J Athl Train. 2014; 49(2):245-265

- 3. Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and
- recommendations for injury prevention initiatives. J Athl Train. 2007; 42(2):311-319
- 4. Laker SR. Epidemiology of concussion and mild traumatic brain injury. American Academy of
- 224 Physical and Med. 2011; 3:354-358
- 5. Daneshvar DH, Nowinski CJ, et al. The epidemiology of sport-related concussion. *Clin Sports Med.* 2011; 30:1-17
- 6. Hunt T, Amato HK. Chiari malformation in a collegiate volleyball player. *Athl Therapy Today*.
 2007; 12(2):12-15.
- 229 7. Nakamura M, Chiba K, Nishizawa T, et al. Retrospective study of surgery-related outcomes in
- 230 patients with syringomyeria associated with Chiari I malformation: clinical significance of
- changes in the size and localization of syrinx on pain relief. J Neurosurg (Spine 3) 2004;
- 232 100:241-244.
- 8. Quon JL. Multimodal evaluation of cerebrospinal fluid (CSF) dynamics following extradural
- decompression for Chiari I malformation. *ProQuest.* 2015; 1-41.
- 235 9. Meehan WP, Jordaan M, Prabhu SP, et al. Risk of athletes with Chiari malformations suffering
- catastrophic injuries during sports participation is low. *Clin J Sports Med.* 2015; 25(2):133-137.
- 10. Harrell BR, Barootes BG. The type I Chiari malformation in a previously asymptomatic
- college athlete: addressing the issue of the return to athletic participation. *Clin J Sport Med.*
- 239 2010; 20(3):215-217.
- 11. Miele VJ, Bailes JE, Martin NA. Participation in contact or collision sports in athletes with
- 241 epilepsy, genetic risk factors, structural brain lesions, or history of craniotomy. *Neurosurg focus*.
- 242 2006; 21(4):1-8.

- 12. Akakin A, Yilmaz B, Eksi MS, et al. Treatment of syringomyelia due to Chiari type I
- malformation with syringe-subarachnoid-peritoneal shunt. *J Korean Neurosur Soc*. 2015;
 57(4):311-313.
- 13. Caldarelli M, Novegno F, Vassimi L, et al. The role of limited posterior fossa craniectomy in
- the surgical treatment of Chiari malformation type I: experience with a pediatric series. J
- 248 Neurosurg. 2007; 106:187-195.
- 249 14. Krischen MP, Illes J. Ethical implications of an incidentally discovered asymptomatic Chiari
- 250 malformation in competitive athlete. *Continuum*. 2014; 20(6):1683-1687.
- 251 15. Starnes TA, Kristy BS, Smith KB, Smith MS. Neurologic symptoms in a football players
- with Chiari I malformation: a case review. *Athletic Training & Sports Health Care*. 2015;7:214253 216.
- 16. Callaway GH, O'Brien SJ, Tehran AM. Chiari I malformation and spinal cord injury: cause
- for concern in contact athletes? *Med Sci Sports Ex.* 1996; 28:1218-1220.
- 256 17. Schuster JM, Zhang F, Norcell DC, et al. Persistent/recurrent syringomyelia after Chiari
- 257 decompression natural history and management strategies: a systematic review. *Evid Based*
- 258 Spine Care J. 2013; 4:116-125.
- 259 18. Ellis JL, Gottlieb JE. Return-to-play decisions after cervical spine injuries. Curr Sports Med
- 260 *Rep.* 2007; 6:56-61.