

11-2010

## ACL reconstructed patients with a BPTB graft present an impaired vastus lateralis neuromuscular response during high intensity running


Kostas Patras  
*University of Ioannina*

Giorgos Ziogas  
*University of Ioannina*

Stavros Ristanis  
*University of Ioannina*

Elias Tsepis  
*Technological Educational Institution of Patras at Aigion*

Follow this and additional works at: <https://digitalcommons.unomaha.edu/biomechanicsarticles>

Nikolaos Stergiou  
 [nstergiou@unomaha.edu](mailto:nstergiou@unomaha.edu)

Please take our feedback survey at: [https://unomaha.az1.qualtrics.com/jfe/form/](https://unomaha.az1.qualtrics.com/jfe/form/SV_8cchtFmpDyGfBLE)

[SV\\_8cchtFmpDyGfBLE](https://unomaha.az1.qualtrics.com/jfe/form/SV_8cchtFmpDyGfBLE)  
See next page for additional authors

### Recommended Citation

Patras, Kostas; Ziogas, Giorgos; Ristanis, Stavros; Tsepis, Elias; Stergiou, Nikolaos; and Georgoulis, Anastasios D., "ACL reconstructed patients with a BPTB graft present an impaired vastus lateralis neuromuscular response during high intensity running" (2010). *Journal Articles*. 86.  
<https://digitalcommons.unomaha.edu/biomechanicsarticles/86>

This Article is brought to you for free and open access by the Department of Biomechanics at DigitalCommons@UNO. It has been accepted for inclusion in Journal Articles by an authorized administrator of DigitalCommons@UNO. For more information, please contact [unodigitalcommons@unomaha.edu](mailto:unodigitalcommons@unomaha.edu).

---

**Authors**

Kostas Patras, Giorgos Ziogas, Stavros Ristanis, Elias Tsepis, Nikolaos Stergiou, and Anastasios D. Georgoulis

1 **ACL RECONSTRUCTED PATIENTS WITH A BPTB GRAFT PRESENT AN IMPAIRED**  
2 **VASTUS LATERALIS NEUROMUSCULAR RESPONSE DURING HIGH INTENSITY**  
3 **RUNNING.**

4 Kostas Patras<sup>1</sup>, Giorgos Ziogas<sup>1</sup>, Stavros Ristanis<sup>1</sup>, Elias Tsepis<sup>1,2</sup>, Nicholas Stergiou<sup>1,3</sup>, Anastasios D.  
5 Georgoulis<sup>1</sup>

6 <sup>1</sup>Orthopaedic Sports Medicine Center, Department of Orthopaedic Surgery, University of Ioannina,  
7 Greece

8 <sup>2</sup>Physical Therapy School, Technological Educational Institution of Patra at Aigion, Greece

9 <sup>3</sup>Nebraska Biomechanics Core Facility, College of Education, University of Nebraska at Omaha and  
10 College of Public Health, University of Nebraska Medical Center, USA

11

12

13

14

15 **Address for Correspondence:**

16 Anastasios D. Georgoulis, M.D

17 Georgiou Papandreou 2

18 Ioannina 45444, GREECE

19 Tel/Fax: +302651064980

20 P.O BOX 1330

21 E-mail: georgoulis@osmci.gr

22

23 Word count

24 ABSTRACT: 232

25 MANUSCRIPT: 3195

26

27

28

## 1 **ABSTRACT**

2 The purpose of the present study was to investigate whether the electromyographic response of the  
3 vastus lateralis (VL) muscle in the anterior cruciate ligament (ACL) reconstructed leg is similar to that  
4 of the intact contralateral leg and healthy controls, during moderate and high intensity running.  
5 Fourteen bone-patellar tendon-bone (BPTB) ACL reconstructed amateur soccer players and fourteen  
6 healthy control amateur soccer players volunteered to participate in the study. Electromyographic  
7 (EMG) traces from the vastus lateralis (VL) muscle were collected bilaterally, as athletes ran on a  
8 treadmill for 10 minutes on separate occasions, at moderate and high intensity. The dependent variable  
9 examined was the EMG amplitude during stance. During the moderate intensity running, EMG  
10 amplitude of the VL did not increase with time for any of the tested legs. During the high intensity  
11 running, the EMG amplitude of the VL increased significantly with time for the intact ( $F=6.747$ ,  
12  $p=0.001$ ) and the control leg ( $F=4.258$ ,  $p=0.008$ ), but remained unchanged for the ACL reconstructed  
13 leg. During moderate intensity running, there was no difference in the neuromuscular response of the  
14 VL in the reconstructed leg compared to the intact and control leg. High intensity running resulted in  
15 an impaired neuromuscular response of the VL in the reconstructed leg compared to the intact and  
16 control leg. It seems that potential impairments of the neuromuscular response after ACL  
17 reconstruction should be tested under high rather than moderate intensity efforts.

18

19 **Key words:** telemetric electromyography, running, ACL reconstruction, exercise intensity

20

21

## 1 INTRODUCTION

2 Anterior cruciate ligament (ACL) function is closely associated with optimal activation of the  
3 muscles surrounding the knee joint [1, 2]. Rupture of the ACL leads to alterations in muscle  
4 recruitment patterns [3, 4]. It has been demonstrated that reconstruction of the ACL re-establishes  
5 muscle activation levels, as assessed by surface electromyography (EMG) amplitude, towards  
6 normative values during moderate intensity activities [5-7]. However, moderate intensity exercises do  
7 not resemble situations that are encountered during participation to sports activities.

8 One feature of the neuromuscular response after ACL reconstruction that has not been  
9 investigated by previous studies is the progression of EMG activation levels with time during the  
10 exercise. It is well established that EMG amplitude time course is greatly influenced by exercise  
11 intensity [8, 9, 10]. During moderate intensity, EMG amplitude does not increase with time [9, 11, 12],  
12 while during high intensity, it does [9, 11-13]. The increasing EMG amplitude with time, during high  
13 intensity exercise is considered to reflect the physiological response that is required for the increasing  
14 metabolic demands of the exercise [8, 9, 10].

15 Previous EMG studies reported that ACL reconstructed (ACLR) subjects exhibit neuromuscular  
16 alterations, such as selective muscle fiber atrophy in the involved vastus lateralis (VL) [14-16], altered  
17 motor unit activation of the quadriceps following surgery and subsequent retraining [17] and loss of  
18 joint afferent information which may lead to suboptimal muscle activation [1, 18]. These  
19 neuromuscular alterations may therefore impair the proper physiological response required at high  
20 exercise intensities. Thus, it is possible that the VL muscle of the operated leg could exhibit impaired  
21 neuromuscular response resulting in inability to naturally increase the EMG amplitude with time  
22 during high intensity exercise.

23 Therefore, the purpose of the present study was to investigate the effect of ACL reconstruction  
24 with BPTB graft on VL activation levels with time during moderate and high intensity running. It is  
25 well established that exercise intensity is best assigned relative to the lactate threshold and  $VO_2$ max [8,  
26 19]. We hypothesized that (a) during moderate intensity exercise the EMG amplitude of the VL will  
27 not increase with time for any of the ACLR, intact and control legs, (b) during high intensity exercise

1 the EMG amplitude of the VL will increase with time for the control and intact leg but not for the  
2 ACLR leg.

3

#### 4 **METHODS**

5 Two groups of athletes participated in the study. Fourteen amateur male soccer players [mean  
6 (SD) age, body weight and height, 24.8 (5.3) years, 77.3 (7.5) kg and 177 (5.3) cm] with ACL-  
7 reconstructed knees and fourteen healthy amateur male soccer layers who had never suffered of any  
8 kind of orthopaedic or neurological condition [mean (SD) age, body weight and height, 21.7 (4.4)  
9 years, 72.2 (8.3) kg and 180 (9.0) cm]. The operated athletes had undergone ACL reconstruction with  
10 bone-patella tendon-bone (BPTB) graft, 18.5 (4.3) months before testing. ACL reconstruction was  
11 performed sub-acutely within 6 months after the injury from the same surgeon (A.G). All subjects had  
12 a unilateral ACL tear confirmed by MRI and arthroscopy.

13 All operated athletes underwent the same rehabilitation protocol, starting from the first post  
14 operative day with the use of passive exercises. Return to sports was permitted 6 months after  
15 reconstruction provided that the athletes had regained stability and full functional strength, according  
16 to well accepted criteria [20]. All subjects agreed with the testing protocol and gave their consent to  
17 participate in accordance with the Institutional Review Board policies of our Medical School. Prior to  
18 any data collection, a clinical evaluation was performed in all athletes by the same clinician. During  
19 this evaluation, the Tegner and Lysholm scores were obtained, while anterior tibial translation was  
20 evaluated using the KT-1000 knee arthrometer (MEDmetric Corp., San Diego, California) [21].

21 The athletes reported to the laboratory on three different occasions, separated by 48 hours,  
22 within a two weeks period. For their first visit to the laboratory, athletes performed an incremental  
23 treadmill (Technogym Runrace 1200, Italy) running test to volitional exhaustion with 3 minute-stages,  
24 to determine  $VO_2$ max and lactate threshold (LT) [22]. A computerized system was used for all  
25 metabolic measurements (CPX Ultima, Medical Graphics, St Maul, MN, USA). At the end of each  
26 stage, capillary blood samples were collected and analyzed for lactate (Accutrend, Roche Diagnostics,  
27 Germany). Prior to each test, all analyzers were calibrated according to the manufacturer instructions.  
28 Attainment of  $VO_2$ max was verified according to criteria established by the American College of

1 Sports Medicine [22]. Lactate threshold was determined according to Cheng et al [23]. The high  
2 intensity running was set at 40% of the difference between  $VO_2$ max and lactate threshold (40%D) and  
3 the moderate intensity running was set at 80% of the lactate threshold [19].

4 In each of the two subsequent visits to the laboratory, athletes were required to perform a 10-  
5 minute run at the pre-selected intensities. We tested one intensity at each visit and the order was  
6 randomly assigned for every athlete. During running, EMG data were collected for 15 seconds at the  
7 3<sup>rd</sup>, 5<sup>th</sup>, 7<sup>th</sup> and 10<sup>th</sup> minute. Gas exchange data were recorded simultaneously breath-by-breath, heart  
8 rate was measured throughout the test and blood lactate was measured prior to running and  
9 immediately after termination of exercise.

10 EMG traces were obtained from the VL muscle bilaterally using bipolar, circular, pre-amplified,  
11 pre-gelled Ag/AgCl electrodes with 10 mm diameter and fixed inter-electrode spacing of 20 mm  
12 (Noraxon Inc, Scottsdale, AZ, USA). EMG data were recorded with a wireless 8-channel EMG system  
13 (Telemetry 2400T, Noraxon Inc, Scottsdale, AZ, USA) and displayed real-time on a personal computer  
14 using dedicated software (MyoResearchXP, Noraxon Inc, Scottsdale, AZ, USA). The surface of the  
15 skin was prepared by shaving hair, rubbing it with abrasive paper and cleaning it with alcohol. The  
16 electrodes were fixed longitudinally over the muscle belly. Electrodes were placed at the antero-lateral  
17 muscle bulge at 2/3 of the proximo-distal thigh length [24, 25]. The visually largest area of muscle  
18 belly was selected using a contraction against manual resistance. The ground electrode was placed on  
19 lateral femoral condyle of the right leg. Electrodes and cables were secured with surgical tape, in  
20 order to avoid any interference with the running pattern of the subjects.

21 Footswitches (Noraxon Inc, Scottsdale, AZ, USA) placed under the heel and big toes of both  
22 legs were used to denote heel-strike and toe-off. EMG was acquired at a sampling rate of 1500 Hz.  
23 The raw EMG was measured in a band of 10 to 500 Hz, full-wave rectified, high pass filtered (cut-off  
24 frequency at 20 Hz) with an 8<sup>th</sup> order Butterworth filter to remove movement artifacts and smoothed  
25 with a 100 ms RMS algorithm.. Values from 20 strides were averaged to calculate the mean peak  
26 amplitude during stance for each of the four time intervals.

27 Based on our hypotheses, the dependent variable examined in the present study was the mean  
28 peak EMG amplitude during the stance phase. A 2-way fully repeated ANOVA within the control

1 group, with time (four levels) and leg (two levels) as within-subjects factors, revealed no time\*leg  
2 interactions for the EMG amplitude for either the moderate or high intensity running. Thus, the left leg  
3 was selected as the control leg.

4 The time course of the EMG amplitude during running was compared between the reconstructed  
5 and intact leg with a 2-way fully repeated ANOVA, with time (four levels) and leg (two levels) as  
6 within-subjects factors. Similarly, the time course of the EMG amplitude during running was  
7 compared between the reconstructed and the control leg and between the intact and control leg, with 2-  
8 way mixed ANOVAs, with time (four levels) as within- and group (two levels) as between-subject  
9 factor. Significant main effects and interactions were investigated with Tukey's post hoc analyses. The  
10 level of significance was set at  $\alpha=0.05$ .

11

## 12 **RESULTS**

13 At the time of data collection no clinical evidence of knee pain and effusion was found in the  
14 ACL reconstructed athletes. All athletes in the ACL-reconstructed group were satisfied with the  
15 outcome of the surgery and resumed their pre-injury level of sports participation. Negative Lachman  
16 and pivot-shift tests indicated that the knee joint stability was regained clinically for all ACL  
17 reconstructed athletes. For the athletes with ACL reconstruction, the median Lysholm score was 95  
18 (range 94-100) and the Tegner score was 8 (range 7-9) at the time of examination. KT-1000 results  
19 revealed that the mean difference between the anterior tibial translation of the reconstructed and intact  
20 sides in the ACL reconstructed group was 1.6 mm (range 1 to 2 mm) for the 134N test and 1.8 mm  
21 (range 1-2 mm) for the maximum manual test, respectively.

22 During the moderate intensity running, EMG amplitude did not increase with time for both the  
23 intact and ACL reconstructed leg ( $F=0.477$ ,  $p=0.7$ ). Similarly EMG amplitude did not increase with  
24 time for both the control and the reconstructed leg ( $F=0.838$ ,  $p=0.477$ ). Finally, EMG amplitude  
25 remained unchanged with time for both the control and intact leg ( $F=0.782$ ,  $p=0.507$ ).

26 During the high intensity running, the EMG amplitude of the VL increased significantly with  
27 time for the intact but not for the reconstructed leg (time\*leg interaction:  $F=6.747$ ,  $p=0.001$ ,  
28 power=0.89). Similarly, the EMG amplitude of the VL increased significantly with time for the



1 control but not for the reconstructed leg (time\*group interaction:  $F=4.258$ ,  $p=0.008$ , power=0.85).  
2 Finally, the EMG amplitude of the VL increased significantly with time for both the control and intact  
3 leg (main effect of time:  $F=11.28$ ,  $p<0.001$ , power=0.99) with no difference between legs (time\*group  
4 interaction:  $F=1.23$ ,  $p=0.303$ ).

5 All effects for the moderate and high intensity running are presented in Table 1.

6 **INSERT TABLE 1 ABOUT HERE**

7

## 8 **DISCUSSION**

9 The purpose of the present study was to investigate the effect of ACL reconstruction with BPTB  
10 graft on VL activation levels with time, during moderate and high intensity running. We hypothesized  
11 that that (a) during moderate intensity running, the EMG amplitude of the VL will not increase with  
12 time for any of the ACLR, intact and control legs, (b) during high intensity running, the EMG  
13 amplitude of the VL will increase with time for the control and intact leg but not for the ACLR leg.

14 Our results indicated that during 10 minutes of moderate intensity running, the EMG amplitude  
15 of vastus lateralis remained unchanged with time for each of the control, intact and reconstructed legs.  
16 Thus, our results supported our first hypothesis. This is in agreement with previous studies indicating  
17 that in healthy individuals performing moderate intensity exercise, EMG amplitude of the exercising  
18 musculature remains unchanged with time [9, 11, 12]. Thus, our results verify that under low demand  
19 activities there are no differences between the reconstructed and either the intact or control leg [5-7].

20 A major finding of the present investigation was that during 10 minutes of high intensity  
21 running, EMG amplitude of the VL increased with time for the control and intact leg but not the  
22 reconstructed leg. Our results confirmed our second hypothesis. Previous studies have shown that  
23 during high intensity exercise, there is a need by the neuromuscular system to compensate for the  
24 accumulating physiological stress [8, 9]. This is accomplished by enhanced activation of the  
25 exercising muscles as exercise progresses and is detected as increased EMG amplitude with time [9,  
26 11-13]. The fact that our two separate groups of athletes responded by increasing EMG amplitude with  
27 time is in good agreement with the above studies. However, the results of our study revealed that  
28 during high intensity exercise the VL muscle of the reconstructed leg failed to respond in a similar

1 manner and showed no increase in EMG amplitude with time compared to either the intact or the  
2 control leg (FIGURE 1). This behaviour may indicate that discrepancies in the neuromuscular  
3 response of the reconstructed leg may be revealed under high intensity sustained athletic efforts.

#### 4 **INSERT FIGURE 1 ABOUT HERE**

5 Several explanations can be given for the lack of increase in EMG amplitude with time for the  
6 VL of the reconstructed leg. These may include selective muscle fiber atrophy in the involved  
7 quadriceps [14-16], altered motor unit activation following surgery and subsequent retraining [17] and  
8 loss of joint afferent information which may lead to suboptimal muscle fiber activation [18]. These  
9 neuromuscular alterations following ACL reconstruction may be responsible for the lack of increase in  
10 the EMG amplitude with time at high intensities, where there is a need for neuromuscular  
11 compensation of the accumulating physiological stress [8, 9, 11-13].

12 To the best of our knowledge this is the first study that investigated EMG activation levels with  
13 time during running in ACL reconstructed and healthy athletes. Previous studies on ACL  
14 reconstructed athletes have compared EMG levels under moderate intensity activities but no study has  
15 investigated the progression of EMG levels with time [5-7]. Furthermore, studies on healthy subjects  
16 have investigated neuromuscular response with time but only during high intensity cycling [11, 12,  
17 13]. Our approach enabled us to extend our findings to intense running which represents a highly  
18 functional activity for the ACL reconstructed athlete. Furthermore, using preliminary  $VO_2$ max and  
19 lactate threshold testing, we assigned moderate and high intensity exercise according to individual  
20 fitness levels. Thus, all subjects exercised under identical controlled conditions.

21 In the present study, we recruited athletes with BPTB autograft. Thus, it is unknown if a similar  
22 response pattern is observed in athletes with a different graft, such as hamstrings. EMG signal  
23 capturing, recording and processing was performed according to established guidelines [27, 28]. We  
24 examined EMG activity developed solely during the stance period, thereby reducing to some extent  
25 the role of the signal non-stationarities [10]. Furthermore, the activity of many successive steps was  
26 averaged providing a reasonable estimation of peak EMG amplitude. Normalisation of EMG data (for  
27 example to maximum voluntary contraction) was not performed due to the additional error introduced  
28 by this process and the fact that our study design involved repeated measures, thereby overcoming

1 influences of electrode positioning and inter-electrode distance on the signal value [26]. Additionally,  
2 we incorporated 2 different control conditions (the intact leg of the reconstructed athletes and the  
3 control leg of a separate group of athletes) to ensure the existence of differences in our dependent  
4 variable. We assumed that because the same instrumentation was used for all subjects, the level of  
5 measurement noise would be consistent for all subjects and that any differences could be attributed to  
6 changes within the system itself.

7

## 8 **Practical applications**

- 9 • Moderate intensity activities may not reveal differences in the neuromuscular response  
10 between the reconstructed and either the intact or the control leg.
- 11 • High intensity sustained efforts are more likely to expose neuromuscular defects of the ACL  
12 reconstructed knee.
- 13 • Endurance type activity may represent a more functional approach to assess neuromuscular  
14 response under fatiguing conditions in ACL reconstructed athletes.

15

## 16 **ACKNOWLEDGEMENTS**

17 This research project (PENED03) is co-financed by E.U.-European Social Fund (75%) and the  
18 Greek Ministry of Development-GSRT (25%).

19

## 20 **REFERENCES**

- 21 1. Johansson H, Sjolander P, Sjoka P. A sensory role for the cruciate ligaments. *Clin Orthop*  
22 *Relat Res* 1991; 268:161-178.
- 23 2. Dyhre Poulsen P, Krosgaard MR. Muscular reflexes elicited by electrical stimulation of the  
24 anterior cruciate ligament in humans. *J Appl Physiol* 2000; 89(6):2191-2195.
- 25 3. Limbird TJ, Shiavi R, Frazer M, et al. EMG profiles of knee joint musculature during  
26 walking: changes induced by anterior cruciate ligament deficiency. *J Orthop Res* 1988;  
27 6(5):630-638.

- 1 4. Van Lent ME, Drost MR, vd Wildenberg FA. EMG profiles of ACL-deficient patients during  
2 walking: the influence of mild fatigue. *Int J Sports Med* 1994; 15(8):508-514.
- 3 5. Bulgheroni P, Bulgheroni MV, Andrini L, et al. Gait patterns after anterior cruciate ligament  
4 reconstruction. *Knee Surg Sports Traumatol Arthrosc* 1997; 5(1):14-21.
- 5 6. Knoll Z, Kiss RM, Kocsis L. Gait adaptation in ACL deficient subjects before and after  
6 anterior ctuciate ligament reconstruction surgery. *J Electromyogr Kinesiol* 2004; 14(3):287-  
7 294.
- 8 7. Lewek M, Rudolph K, Axe M, et al. The effect of insufficient quadriceps strength on gait after  
9 anterior cruciate ligament reconstruction. *Clin Biomech* 2002; 17(1):56-63.
- 10 8. Gaesser GA, Poole DC. The slow component of oxygen uptake kinetics in humans. *Exerc*  
11 *Sport Sci Rev* 1996; 24:35-71.
- 12 9. Jones AM, Pringle JSM, Carter H. Influence of muscle fiber type and motor unit recruitment  
13 on VO<sub>2</sub> kinetics, Chapter 11, in *Oxygen uptake kinetics in sport, exercise and medicine*. Jones  
14 AM, Poole DC, eds London. Routledge; 2005.
- 15 10. De Luca CJ. The use of surface electromyography in biomechanics. *J Appl Biomech* 1997;  
16 13:135-163.
- 17 11. Shinohara M, Moritani T. Increase in neuromuscular activity and oxygen uptake during heavy  
18 exercise. *Ann Physiol Antropol* 1992; 11:257-262.
- 19 12. Saunders MJ, Evans EM, Arngrimsson SA, et al. Muscle activation and the slow component  
20 rise in oxygen uptake during cycling. *Med Sci Sports Exerc* 2000; 32(12):2040-2045.
- 21 13. Sabapathy S, Schneider DA, Morris NR. The VO<sub>2</sub> slow component: relationship between  
22 plasma ammonia and EMG activity. *Med Sci Sports Exerc* 2005; 37(9):1502-1509.
- 23 14. Bryant AL, Kelly J, Hohmann E. Neuromuscular adaptations and correlates of knee  
24 functionality following ACL reconstruction. *J Orthop Res* 2008; 26(1):126-135.
- 25 15. McHugh MP, Tyler TF, Nicholas SJ, et al. Electromyographic analysis of quadriceps fatigue  
26 after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 2001; 31(1):25-32.

- 1 16. Snyder-Mackler L, Binder-Macleod SA, Williams PR. Fatigability of human quadriceps  
2 femoris muscle following anterior cruciate ligament reconstruction. *Med Sci Sports Exerc*  
3 1993; 25(7):783-789.
- 4 17. Drechsler WI, Cramp MC, Scott OM. Changes in muscle strength and EMG median  
5 frequency after anterior cruciate ligament reconstruction. *Eur J Appl Physiol* 2006; 98(6):613-  
6 623.
- 7 18. Konishi Y, Fukubayashi T, Takeshita D. Mechanism of quadriceps femoris muscle weakness  
8 in patients with anterior cruciate ligament reconstruction. *Scand J Med Sci Sports* 2002;  
9 12(6):371-375.
- 10 19. Carter H, Pringle JSM, Jones AM, et al. Oxygen uptake kinetics during treadmill running  
11 across exercise intensity domains. *Eur J Appl Physiol* 2002; 86(4):347-354.
- 12 20. DeLee J, Drez D, Miller M, eds. DeLee & Drez's orthopaedic sports medicine: principles and  
13 clinical practise 2<sup>nd</sup> ed. Philadelphia, PA. Saunders. 2003.
- 14 21. Daniel DM, Malcom LL, Losse G, et al. Instrumented measurement of anterior laxity of the  
15 knee. *J Bone Joint Surg Am* 1985; 67(5):720-6.
- 16 22. Whaley MH, Brubaker PH, Otto RM, eds. ACSM's Guidelines for Exercise Testing and  
17 Prescription, 7<sup>th</sup> ed. Philadelphia, PA. Lippincott Williams & Wilkins. 2005.
- 18 23. Cheng B, Kuipers A, Snuder C et al. A new approach for the determination of ventilatory and  
19 lactate thresholds. *Int J Sports Med* 1992; 13(7):518-22.
- 20 24. Freriks B, Hermens H, Disselhorst-Klug C, et al. The recommendations for sensors and sensor  
21 placement procedures for surface electromyography, in European recommendations for  
22 surface electromyography, Hermens HJ, ed.. Enschede. Roessingh Research and  
23 Development. 1999.
- 24 25. Merletti R and Hermens HJ. Detection and conditioning of the surface EMG signal, in:  
25 Electromyography: Physiology, Engineering and Non-Invasive Applications, Merletti R and  
26 Parker P, eds. New Jersey. Wiley-IEEE, 2004.

1 26. Rahnama N, Lees A and Reilly T. Electromyography of selected lower-limb muscles fatigued  
2 by exercise at the intensity of soccer match-play. *J Electromyogr Kinesiol* 2006; 16(3): 257-  
3 263.  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

1 **FIGURE LEGENDS**

2 **FIGURE 1:** (a) EMG amplitude over time for the intact (n=14) and ACL reconstructed leg (n=14)  
3 during the high intensity running. EMG amplitude increased compared to initial values only for the  
4 control leg. \* indicates significantly higher than initial value. Vertical bars represent  $\pm 95\%$  confidence  
5 intervals. (b) EMG amplitude over time for the control (n=14) and ACL reconstructed leg (n=14)  
6 during the high intensity running. EMG amplitude increased compared to initial values only for the  
7 intact leg. \* indicates significantly higher than initial value. Vertical bars represent  $\pm 95\%$  confidence  
8 intervals.

9