



The Effects of Trichloroethylene on the Germination and Early Development of Radish, Butternut Squash, Corn, and Soybean

Sarah Jaksich, and P. Roxanne Kellar

Department of Biology, University of Nebraska at Omaha, Omaha, NE 68182

ABSTRACT

Trichloroethylene (TCE) is a small molecule used as a metal degreaser and as an extraction solvent for lipids. Historically, TCE has been used to clean missiles and rockets after testing at military sites. However, TCE has decreased in use due to its potential health hazards. TCE is particularly dangerous because it is soluble in water and easily seeps into ground water. As a result, TCE has contaminated groundwater in Mead, Nebraska. The town is near the former Nebraska Ordnance Plant, a military factory specializing in bomb assembly, ammonium nitrate production, and explosives burning. Soil sampling in Mead indicated TCE levels as high as 10,000 µg/L, with most positive sites testing greater than 10 µg/L. There is little information concerning the effects of TCE on local flora or agriculture, most of it focusing on human health. In this study, we evaluated the effects of TCE on germination and early seedling development of radish, butternut squash, corn, and soybean. Seeds were dosed with TCE concentrations ranging from 0 µg/L – 100,000 µg/L and allowed to grow until the first foliage leaf emerged. We observed each developmental landmark, and measured root length. This knowledge will give insight into the effects of TCE on the development of common Nebraska crops and aid in agricultural decision making.

METHODS

RESULTS

INTRODUCTION

REFERENCES

- Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological Profile for Trichloroethylene (Update). U.S. Public Health Service, U.S. Department of Health and Human Services, Atlanta, GA. 1997.
- Farmer, S. (2016, May 25). New wells digging deeper at former NOP. Retrieved January 16, 2019, from https://www.wahoo-ashland-waverly.com/news/new-wells-digging-deeper-at-former-nop/article_4d6e03e4-228e-11e6-9f9d-af36ba203a42.html
- NWK. 2013 TCE and RDX Concentrations [PDF]. (2014). Kansas City: US Army Corps of Engineers.
- Ryu, S. B., Davis, L. C., Selk, K., & Erickson, L. E. (1996). Evaluation of Toxicity of Trichloroethylene for Plants [PDF]. Manhattan KS: Departments of Biochemistry and Chemical Engineering, Kansas State University.
- Schnabel, William & Dietz, Annette & Burken, Joel & Schnoor, Jerald & J. Alvarez, Pedro. (1997). Uptake and transformation of trichloroethylene by edible garden plants. *Water Research*. 31. 816-824.
- Schöftner, P., Watzinger, A., Holznecht, P., Wimmer, B., & Reichenauer, T. G. (2016). Transpiration and metabolisation of TCE by willow plants – a pot experiment. *International Journal of Phytoremediation*, 18(7), 686–692.
- Strycharz, S., & Newman, L. (2009). Use of Native Plants for Remediation of Trichloroethylene: I. Deciduous Trees. *International Journal of Phytoremediation*, 11(2), 150–170.
- U.S. Environmental Protection Agency. Trichloroethylene Health Risk Assessment: Synthesis and Characterization. External Review Draft. EPA/600/P-01/002A. Office of Research and Development, Washington, DC. August 2001.

ACKNOWLEDGEMENTS

We thank the Funding for Undergraduate Scholarly Experiences (FUSE) and the University Committee on Research and Creative Activity (UCRCA) grant awarded by UNO.