

Student Work

4-1-1998

Tallgrass Prairie Remnants of Western Iowa Cemeteries.

Carrie L. Menges-Schaben

Follow this and additional works at: <https://digitalcommons.unomaha.edu/studentwork>

Recommended Citation

Menges-Schaben, Carrie L., "Tallgrass Prairie Remnants of Western Iowa Cemeteries." (1998). *Student Work*. 3338.

<https://digitalcommons.unomaha.edu/studentwork/3338>

This Thesis is brought to you for free and open access by DigitalCommons@UNO. It has been accepted for inclusion in Student Work by an authorized administrator of DigitalCommons@UNO. For more information, please contact unodigitalcommons@unomaha.edu.



Tallgrass Prairie Remnants of Western Iowa Cemeteries

A Thesis

Presented to the

Department of Biology

and the

Faculty of the Graduate College

University of Nebraska

In Partial Fulfillment

Of the Requirements for the Degree

Masters of Arts in Biology

University of Nebraska at Omaha

By

Carrie L. Menges-Schaben

April 1998

UMI Number: EP74940

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI EP74940

Published by ProQuest LLC (2015). Copyright in the Dissertation held by the Author.

Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code



ProQuest LLC.
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106 - 1346

THESIS ACCEPTANCE

Acceptance for the faculty of the Graduate College,
University of Nebraska, in partial fulfillment of the requirements
for the degree Master of Arts, University of Nebraska at Omaha.

Committee

Name	Department/School
<i>David M. Sutherland</i>	<i>Biology</i>
<i>Jeffrey T. Peckham</i>	<i>Geography</i>

Chairperson *Thomas P. Burg*

Date *21 April 1998*

TABLE OF CONTENTS

Abstract.....	ii
Introduction.....	1
Methods.....	8
Results.....	11
Discussion.....	19
Literature Cited.....	22
Appendix.....	28
Table 1.....	29

LISTS OF TABLES AND FIGURES

TABLES

Table 1. Characteristics of cemetery sites.....	10
Table 2. Mean canopy cover and S.E. of all species.....	12
Table 3. Shannon-Weiner Diversity values.....	15

FIGURES

Fig. 1. Location of tallgrass prairie remnant.....	9
Fig. 2. Mean canopy cover of big bluestem.....	17
Fig. 3. Mean canopy cover of smooth brome.....	18

ABSTRACT

Tallgrass prairie remnants situated in eight western Iowa cemeteries were sampled during 1995 and 1996 to compare both the frequency and season of mowing and burning on plant species composition. Dominant, native tallgrass prairie species, such as big bluestem (*Andropogon gerardii*), Indian grass (*Sorghastrum nutans*) and leadplant (*Amorpha canescens*), generally showed a significant increase with burning, whether mowed or not. Mowing, however, adversely affected other species, including flowering spurge (*Euphorbia corollata*) and porcupine grass (*Stipa spartea*), whether burned or not. Non-native species, in particular smooth brome (*Bromus inermis*) increased with mowing and the absence of fire. Overall, my study suggests the importance of selecting the appropriate type or frequency of management in order to maintain the native tallgrass prairie plant diversity. Further, it supports the appropriate use of fire rather than mowing, both to favor native species and to limit the advance of non-native species.

INTRODUCTION

An estimated 90% of the original 58 M ha of tallgrass prairie (*Andropogon-Panicum-Sorghastrum*) (Küchler 1964, 1985) have been lost with extant prairie remnants occurring mostly in small fragments (Klopatek *et al.* 1979). Regional losses vary from 85% west of the Missouri River to as much as 99.9% in Iowa (Samson and Knopf 1994). Losses are largely a consequence of such factors as cultivation, urban expansion, and alteration of natural fire regime that has exposed prairies to invasion by woody plants (Bragg and Hulbert 1976) or non-native, herbaceous species (Lovell *et al.* 1983).

Tallgrass prairie remnants in Iowa are most commonly found either along roadsides or within the boundaries of old cemeteries (Hanes and Hanes 1947, Kerr and White 1981). In these sites, mowing and burning are the principal management options available to land managers. Which option to apply and the manner (e.g. season and frequency) by which to apply it are dictated by various considerations including prairie location, local expertise, management objectives, and the degree of concern for invasive, native woody species and non-native, herbaceous species, such as smooth brome (*Bromus inermis*), Kentucky bluegrass (*Poa pratensis*) and musk thistle (*Carduus nutans*).

Management of most of Iowa's prairie remnants has involved either mulch-mowing (i.e. mowing and not removing the mulch) or haying (i.e. mowing and removing the mulch). In this study, use of the term *mowing* will refer to mulch-mowing while use of the term *haying* will refer to mowing and removing mulch. Mowing and haying have been noted to have different effects in grasslands. Dale (1983), for example, found that grasses showed

better, initial growth on plots where mulch was removed than where mulch remained, probably a response to a warmer soil and more adequate moisture. By mid-July, however, those plots with mulch showed greater growth, probably due to moisture retained in soil by the mulch cover. Over time, however, haying may result in a decline in ecosystem productivity (Ehrenreich and Aikman 1963). Further, repeatedly mowing or haying tallgrass prairie appears to favor some cool-season species, particularly the aggressive eurasian grass, smooth brome, which has the potential to replace native species (Boettcher and Bragg 1989).

More recently, prescribed burning has been used in prairie management, since fire was historically a natural component of the tallgrass prairie ecosystem (Pyne 1982) and since fire can also be a more cost-effective means by which to manage grasslands (Engle 1988). Fire's role in tallgrass prairie is varied (Bragg 1995). Effects of burning include (1) woody plant suppression (Bragg and Hulbert 1976), (2) invigoration of prairie plants and their reproduction through litter removal and increased soil temperature (Hulbert 1969, 1986), and (3) control of invasive, herbaceous species (Lovell *et al.* 1983). In the absence of fire, not only do native woody species invade but so do non-native species such as Kentucky bluegrass and smooth brome (Ross and Vanderpoel 1991).

Comparisons between burning and mowing.- A comparison between burning, a natural process, and mowing, a human process, is necessary in order to assess the degree to which one may be used as a surrogate for the other. In general, mowing and burning are similar in that both (1) may maintain a high diversity of grassland species while controlling woody plant

invasion (Mooberry 1984, Solecki and Toney 1986, Fitzgerald and Tanner 1992), (2) may affect either the amount or distribution of standing dead and litter, and (3) typically require planning decisions such as the time and frequency of application and long-term management goals. Some differences, however, are apparent when comparing burning with either haying or mulch-mowing. Haying is more similar to burning than mulch-mowing because both haying and burning remove plant matter, thereby minimizing litter accumulation. Differences, however include that haying does not remove all litter although it does generally remove nutrients from the system.

While there are some similarities, the differences between burning and mowing are more substantial. Most importantly, fire stimulates the tallgrass prairie plant community more than does mowing, although neither all species nor all prairies respond to the same degree (Hadley 1970, Pemble *et al.* 1981). In addition, the blackened surface resulting from burning causes soil temperatures to be higher due to higher absorption of incident solar radiation that may be responsible for spring plant growth being advanced by as much as 2-3 weeks (Ehrenreich 1959, Ehrenreich and Aikman 1963, Hulbert 1969). These differences may also affect microbial responses. Yet another difference is that both haying and mulch-mowing leave a more uniform distribution of plant matter than does burning (Morris 1979). In addition, mowing may remove woody plant cover more completely, particularly that part that is above mowing height, whereas burning may only reduce the amount or extent of woody plant cover and often only temporarily (Evans 1983, Fitzgerald and Tanner 1992).

Certain logistical differences also occur between burning and mowing.

For example, mowing is more easily applied at different seasons than is fire and it can also be applied several times each growing season. On the other hand, burning is usually limited to a single treatment each year, depending on the rate of fuel (litter) accumulation. For example, spring-burned tallgrass prairie accumulates enough fuel so that it can be reburned again in the fall of that year (Bragg 1982) whereas slow litter accumulation in short-grass prairie limits fire frequency to 6-10 years (Leenhouts 1997). The season and frequency of application, thus, are among important variables for both mowing and burning (Aldous 1934, McMurphy and Anderson 1965).

Frequency of treatment.- In general, frequent treatment by either mowing or burning is detrimental to the native tallgrass prairie plant community. Frequent mowing (e.g. annually or several times in a single year), for example, results in both a decline in native species diversity and an increase in disturbance species (Boettcher and Bragg 1989) although it also provides better control of woody invaders than does either a single mowing or a single prescribed fire (Evans 1983). Similarly, frequent burning results in both a decline in productivity (Conard 1954) and a change in community composition (McMurphy and Anderson 1965). For example Kucera and Koelling (1964) and Solecki and Toney (1986) found that, in contrast to an unburned area, annual burning resulted in a uniform grass cover with a reduction in forbs, whereas, burning every other year resulted in a greater number of forbs. Burning too infrequently, however, had nearly the same results as not burning. Communities burned at 5-year intervals, for example, differed little from those that were unburned (Kucera and Koelling 1964).

Season of treatment.- Seasonal effects of mowing or burning tallgrass prairie are generally divided into spring, summer, and fall. Spring is the season when C₃ (cool-season) species are actively growing but before growth of the C₄ (warm-season) species that dominate the region. Summer is the time of rapid growth of C₄ species but also the time of most lightning-caused fires (Bragg 1982) whereas fall is the season during which C₄ species are dormant but C₃ species may be active.

The season of treatment greatly affects individual and community-level responses to mowing. Spring (April) mowing tends to favor C₄ species (Hulbert 1969, Launchbaugh and Owensby 1978, Hover and Bragg 1981, Gillen and McNew 1987) and thus may be used to affect some control of introduced C₃ species such as Kentucky bluegrass (Curtis and Partch 1948, Ehrenreich and Aikman 1963), smooth brome (Solecki and Toney 1986) and to a lesser extent non-prairie woody plants (Lovell *et al.* 1983). The greatest diversity of prairie grasses and forbs, however, appears to occur on tallgrass prairies disturbed with summer (May-August) mowing (Launchbaugh and Owensby 1978, Solecki *et al.* 1986, Solecki and Toney 1986) although, annual summer mowing, over the long-term, decreases net productivity (Ehrenreich and Aikman 1963). Dormant-season mowing (October-February), whether periodic or annual, will retain at least some of native tallgrass species (Hayden and Aikman 1949) but generally reduces native plant diversity and leads to invasion by C₃ species such as Kentucky bluegrass and smooth brome (Weaver and Fitzpatrick 1934, Hayden and Aikman 1949, Launchbaugh and Owensby 1978, Boettcher and Bragg 1989,

Gibson *et al.* 1993).

As with mowing, the season of occurrence of burning greatly affects individual and community-level responses. In tallgrass prairies dominated by C₄ species, spring burns more consistently decrease C₃ species, whether non-native species, such as Kentucky bluegrass, or native species, such as porcupine-grass (*Stipa spartea* Trin.), Canada wild rye (*Elymus canadensis* L.), and Scribner dichanthelium [*Dichanthelium oligosanthos* (Schult.) Gould var. *scribnerianum* (Nash) Gould] (Hensel 1923, Robocker and Miller 1955, Ehrenreich 1959, Hadley and Kieckhefer 1963, Anderson 1965, Old 1969). Spring burning also affects the forb community, although, late spring burning reduces canopy cover of all forbs to a greater extent than does earlier burning (McMurphy and Anderson 1965, Towne and Owensby 1984, Hulbert 1988). In contrast to spring burning, summer burns have been shown to be most successful in effecting some degree of woody plant control (Hulbert 1986). Fire in this season, however, may open sites to potential erosion and to invasion by exotic species (Solecki and Toney 1986) but it has also been shown to favor some prairie species such as prairie clover (*Dalea* spp.) and false sunflower (*Heliopsis helianthoides* var. *scabra*) (Bragg 1991). Fall burning, in general, has been found to increase rigid sunflower (*Solidago rigida* L.) (Schwegman and McClain 1985), leadplant (*Amorpha canescens* Pursh) (Towne and Owensby 1984) and white sweet clover (*Melilotus alba* Medic.) (Schwegman and McClain 1985). Annual fall burning may also increase forb diversity (Kline 1986) although it tends to lower C₄ grass production (Hulbert 1986).

Effects of no treatment.- In the absence of some form of management, (e.g. burning or mowing), litter accumulates, net annual primary production decreases, and aggressive plants, including non-native species, may increasingly replace prairie species (Hayden and Aikman 1949, Ehrenreich and Aikman 1963, Hulbert 1986, Knapp and Seastedt 1986, Leach and Givnish 1996). Studies on seasonal fire effects by Bragg (1991), for example, found that in both eastern Nebraska tallgrass prairies and northwestern Iowa loess hills prairies, forb cover declined most in areas that were not burned. Of particular concern, however, is the replacement of native prairie species by non-native exotics such as smooth brome and sweet-clover (Ross and Vanderpoel 1991). Once established, removal of these species, particularly smooth brome, can be difficult at best. For example, neither burning nor mowing was able to control smooth brome that had become established in a reseeded tallgrass prairie (Willson and Stubbendieck 1996). Further, where management may be suitable to control invasive species, it often may not be optimal for maintaining a desired prairie composition (Lovell *et al.* 1983). Over longer periods of time, woody plant encroachment of untreated areas is likely to degrade tallgrass prairie species composition (Tomanek 1948, Gehring and Bragg 1992) or replace tallgrass prairie with forest (Bragg and Hulbert 1976).

Only small stands of historic tallgrass prairie remain, especially in western Iowa, and maintaining the native diversity of these remnants is of particular ecological importance. Thus, there is a need to further identify and understand extant prairie remnants and draw from that knowledge that which will allow us to apply available management options intelligently in the future. Since historic cemeteries are one of the principal locations of western Iowa

tallgrass prairie remnants, this study was initiated to identify the cemetery prairie management regime most likely to be successful in maintaining the greatest amount of native tallgrass prairie diversity.

METHODS

Study Area.- The study was conducted in western Shelby and Crawford counties of western Iowa. In 1994, eight cemetery sites, each containing native, tallgrass prairie, were selected for study based on relatively similar (1) previous management, (2) soil type, (3) topographic location, (4) presence of native tallgrass prairie species, and (5) no known history of inter-seeding or use of herbicides (Fig. 1). Sites selected included (1) three sites that, for many years, had been mulch-mowed three times annually and that were burned once in the spring of 1995 (mowed-burned), (2) three sites that had not been historically mulch-mowed but that were burned in 1995 (unmowed-burned), and (3) two sites that had a history of mowing but that were not burned (mowed-unburned) (Fig. 1). Permission to conduct research was obtained from the local cemetery boards with site management organized through the local county conservation boards.

Soils of the sites were generally well drained, silty, loess soils of the Mollisol or Entisol soil orders (Table 1). Climate of the region is continental, with normal daily highs averaging 30 C (86 F) in July and -12 C (10.9 F) in January. Annual precipitation (based on 1951 to 1980 data) averages 76 cm, with 74% occurring during the growing season (April through September) (National Oceanic and Atmospheric Association 1989).

Data Collection.- At each site, vegetation was evaluated in ten 30 by

Fig. 1. Location of tallgrass prairie remnants of western Iowa cemeteries.




Fig. 1. 1-Manteno Cemetery
 2-Holcomb Cemetery
 3-Washington Township Cemetery, West
 4-Washington Township Cemetery, East
 5-Buckgrove Cemetery
 6-Catholic Cemetery
 7-King Cemetery
 8-Willow Cemetery

Table 1. Characteristics of cemetery sites in Shelby and Crawford Counties, Iowa.

Site No.	Name	Treatment	Size (ha)	Location	Soil	
					Type	Order
1	Buckgrove	Unmowed-Burned	0.8	NE 1/4 of NE 1/4 of Sect. 21	Monona-Ida Silt Loam	Mollisol-Entisol
2	Vail	Unmowed-Burned	4.0	NE 1/4 of NE 1/4 of Sect. 25	Monona Silt Loam	Mollisol
3	Willow	Unmowed-Burned	2.0	SW 1/4 of NW 1/4 of Sect. 22	Colo-Napier-Nodaway Silt Loam	Mollisol-Mollisol-Entisol
4	Manteno	Mowed-Burned	0.4	SW 1/4 of SW 1/4 of Sect. 8	Judson Silty Clay Loam	Mollisol
5	Washington Township-East	Mowed-Burned	1.0	NW 1/4 of NE 1/4 of Sect. 22	Monona-Ida Silt Loam	Mollisol-Entisol
6	Catholic	Mowed-Burned	0.8	NW 1/4 of SE 1/4 of Sect. 18	Marshall Silty Clay Loam	Mollisol
7	Holcomb	Mowed-Unburned	0.8	NW 1/4 of NW 1/4 of Sect. 14	Monona Silt Loam	Mollisol
8	Washington Township-West	Mowed-Unburned	0.6	NW 1/4 of NE 1/4 of Sect. 22	Monona-Ida Silt Loam	Mollisol-Entisol

50 cm quadrats that were randomly located within hilltop locations. Sampling was conducted during both late spring (June) and late summer (September) in both 1995 and 1996 in order to ensure that the greatest number of resident species were detected, particularly those evident in the spring but not so in the fall. Sampling consisted of estimating canopy cover for each species in nine categories: absent, 0-1%, >1-5%, >5-25%, >25-50%, >50-75%, >75-95%, >95-99%, >99% cover (modified from Daubenmire 1959). Midpoint values were used for analysis. Voucher specimens were collected and are located in the University of Nebraska at Omaha Herbarium (OMA). Nomenclature follows the Great Plains Flora Association (1986).

Significant differences among and between treatment, by species, were based on a non-parametric, ranked ANOVA and the Student-Newman-Keuls (SNK) multicomparison tests, (SAS Institute Inc. 1985). Statistical comparisons between sites of Shannon-Weiner Diversity Indices used procedures described in Zar (1984).

RESULTS

Community Response.- Forty plant species were identified during this study of which sixteen were non-native (Table 2). The greatest number of native plants ($S = 297$) was found in one of the mowed-burned treatments and the greatest diversity of native species was found in unmowed-burned treatments ($H' = 1.1716$) (Table 3). In contrast, the lowest number of native plants ($S = 45$) was found in a mowed-unburned site, although, diversity was lowest ($H' = 0.5827$) in a mowed-burned treatment. When compared by treatment, no significant differences were found between grasses, forbs and woody plant cover.

Table 2. Mean canopy cover and Standard Error of all species in cemetery sites. Alphabetic superscripts that differ between treatments indicate a significant difference ($p < 0.05$) between mean values for the species, season and year indicated. tr = <1% cover; S = spring sampling, F=fall sampling. * = Non-native species. n=30 for mowed-burned and unmowed-burned treatments and n=20 for mowed-unburned treatment.

Common Name (<i>Scientific name</i>)	Season-Year of Evaluation	Treatment		
		Mowed Burned	Unmowed Burned	Mowed Unburned
Shannon-Weiner Diversity Index (H')	F95	0.6676	0.4981	0.8113
	F96	0.8351	0.9257	0.8032
Forbs	F96	32±2.5 ^a	44±3.4 ^a	39±3.7 ^a
Grasses	F96	70±2.9 ^a	65 ±3.4 ^a	70±3.9 ^a
Woody Plant	F96	0±0.0 ^a	0 ±0.0 ^a	0±0.0 ^a
Lead Plant (<i>Amorpha canescens</i>)	F95	4±2.2 ^{ab}	8 ±2.9 ^a	tr ^a
	F96	2±0.8 ^a	6±1.9 ^a	6±2.2 ^a
Big Bluestem (<i>Andropogon gerardii</i>)	F95	61±6.0 ^a	62±4.7 ^a	27±7.0 ^b
	F96	29±3.8 ^a	22±2.9 ^a	28±4.9 ^a
Little bluestem (<i>Andropogon scoparius</i>)	F95	0±0.0 ^a	0 ±0.0 ^a	0±0.0 ^a
	F96	1± 1.3 ^b	5±1.9 ^a	tr ^{ab}
Meadow Anemone (<i>Anemone cylindrica</i>)	F95	0±0.0 ^a	0 ±0.0 ^a	0±0.0 ^a
	F96	tr ^a	0 ±0.0 ^a	tr ^a
Common Milkweed (<i>Asclepias syriaca</i>)	F95	1±0.5 ^a	0 ±0.0 ^a	0±0.0 ^a
	F96	0±0.0 ^a	0 ±0.0 ^a	0±0.0 ^a
Whorled Milkweed (<i>Asclepias verticillata</i>)	F95	0±0.0 ^a	0 ±0.0 ^a	0±0.0 ^a
	F96	0±0.0 ^a	0 ±0.0 ^a	0±0.0 ^a
Sideoats Grama (<i>Bouteloua curtipendula</i>)	F95	20±4.9 ^a	1 ±0.5 ^b	12±4.3 ^{ab}
	F96	8±2.1 ^a	5 ±1.5 ^a	4±1.9 ^a
Smooth Brome* (<i>Bromus inermis</i>)	F95	6±2.1 ^a	2±0.8 ^b	23±4.9 ^a
	F96	6±1.6 ^a	2±1.0 ^b	8±2.2 ^a
Field Bindweed* (<i>Convolvulus arvensis</i>)	F95	0±0.0 ^a	0 ±0.0 ^a	0±0.0 ^a
	F96	1±0.5 ^a	1±0.5 ^a	0 ±0.0 ^a
Finger Coreopsis (<i>Coreopsis palmata</i>)	F95	0±0.0 ^a	tr ^a	0±0.0 ^a
	F96	tr ^a	1±1.3 ^a	tr ^a
Crown Vetch* (<i>Coronilla varia</i>)	F95	0±0.0 ^a	0 ±0.0 ^a	0±0.0 ^a
	F96	0±0.0 ^b	0±0.0 ^b	3±1.4 ^a

Table 2. Mean canopy cover \pm S.E. of all species (continued).

Common Name (<i>Scientific name</i>)	Season-Year of Evaluation	Mowed Burned	Unmowed Burned	Mowed Unburned
Orchard Grass* (<i>Dactylis glomerata</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
White Prairie Clover (<i>Dalea candida</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	2 \pm 1.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Prairie Larkspur (<i>Delphinium virescens</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Scribner Dichanthelium (<i>Dicanthelium oligosanthes</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^b	2 \pm 0.8 ^a	0 \pm 0.0 ^b
Scouring Rush (<i>Equisetum hyemale</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	tr ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Daisy Fleabane (<i>Erigeron strigosus</i>)	F95	2 \pm 1.0 ^a	0 \pm 0.0 ^a	1 \pm 0.8 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	tr ^a
Flowering Spurge (<i>Euphorbia corollata</i>)	F95	0 \pm 0.0 ^b	7 \pm 2.7 ^a	tr ^b
	F96	0 \pm 0.0 ^c	8 \pm 1.9 ^a	1 \pm 0.2 ^b
White Avens (<i>Geum canadense</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
False Sunflower (<i>Heliopsis helianthoides</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	tr ^a	0 \pm 0.0 ^a	1 \pm 0.8 ^a
Wild Lettuce* (<i>Lactuca canadensis</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	tr ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Black Medic* (<i>Medicago lupulina</i>)	F95	0 \pm 0.0 ^a	4 \pm 2.2 ^a	2 \pm 1.2 ^a
	F96	7 \pm 1.6 ^a	5 \pm 1.6 ^{ab}	1 \pm 0.3 ^b
White Sweet Clover* (<i>Melilotus alba</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Yellow Sweet Clover* (<i>Melilotus officinalis</i>)	F95	3 \pm 1.7 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	1 \pm 0.5 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Yellow Wood Sorrel* (<i>Oxalis stricta</i>)	F95	0 \pm 0.0 ^a	tr ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Prairie Phlox (<i>Phlox pilosa</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a

Table 2. Mean canopy cover \pm S.E. of all species (continued).

Common Name (<i>Scientific name</i>)	Season-Year of Evaluation	Mowed Burned	Unmowed Burned	Mowed Unburned
Timothy* (<i>Phleum pratense</i>)	F95	1 \pm 0.5 ^a	0 \pm 0.0 ^a	4 \pm 2.9 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Blackseed Plantain* (<i>Plantago rugelii</i>)	F95	6 \pm 2.4 ^a	0 \pm 0.0 ^b	0 \pm 0.0 ^b
	F96	1 \pm 0.7 ^a	0 \pm 0.0 ^b	0 \pm 0.0 ^b
Kentucky Bluegrass* (<i>Poa pratensis</i>)	F95	0 \pm 0.0 ^b	0 \pm 0.0 ^b	21 \pm 6.4 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	tr ^a
Bur Oak (<i>Quercus macrocarpa</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Grayhead				
Prairie Coneflower (<i>Ratibida pinnata</i>)	F95	2 \pm 1.3 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	1 \pm 0.7 ^a	tr ^a
Prairie Wild Rose (<i>Rosa arkansana</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	1 \pm 0.7 ^a	0 \pm 0.0 ^a
Black Eyed Susan (<i>Rudbeckia hirta</i>)	F95	1 \pm 0.7 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Yellow Foxtail* (<i>Setaria glauca</i>)	F95	tr ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	4 \pm 1.1 ^a	0 \pm 0.0 ^b	4 \pm 1.4 ^a
Prairie Goldenrod (<i>Solidago missouriensis</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
	F96	tr ^a	2 \pm 0.8 ^a	0 \pm 0.0 ^a
Indian Grass (<i>Sorghastrum nutans</i>)	F95	1 \pm 0.7 ^a	1 \pm 0.5 ^a	0 \pm 0.0 ^a
	F96	25 \pm 3.6 ^a	26 \pm 4.5 ^a	5 \pm 1.5 ^b
Porcupine Grass (<i>Stipa spartea</i>)	F95	0 \pm 0.0 ^a	1 \pm 0.5 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Goats beard* (<i>Tragopogon dubius</i>)	F95	0 \pm 0.0 ^a	0 \pm 0.0 ^a	2.0 \pm 1.9 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a
Red Clover* (<i>Trifolium pratense</i>)	F95	1 \pm 0.7 ^b	1 \pm 0.5 ^b	8 \pm 2.7 ^a
	F96	8 \pm 1.6 ^a	2 \pm 0.8 ^b	2 \pm 1.2 ^b
Blue Vervain (<i>Verbena stricta</i>)	F95	0 \pm 0.0 ^a	1 \pm 0.5 ^a	0 \pm 0.0 ^a
	F96	0 \pm 0.0 ^a	0 \pm 0.0 ^a	0 \pm 0.0 ^a

Table 3. Shannon-Weiner Diversity values (H') based on all species collected in spring and fall of both 1995 and 1996, at eight Iowa cemetery sites. S²H'=variance of sites, s=species richness. Cemetery treatments for sites are as follows: 1-3 =unmowed-burned (UM-B), 4-6 = mowed-burned (M-B), 7-8 = mowed-unburned (M-UB).

Treatment	Site	Native Species		All Species	
		S ² H'	H'	S ² H'	H'
UM-B	1	183	0.9638	183	0.9829
UM-B	2	195	0.7124	195	0.9077
UM-B	3	237	1.1716	237	1.1716
M-B	4	166	0.5827	166	0.8286
M-B	5	293	1.1267	293	1.1267
M-B	6	181	0.6995	181	0.6995
M-UB	7	45	0.8866	45	0.8866
M-UB	8	175	0.9514	175	0.9514

Individual Species Response.-

Burning.- Maximum annual canopy cover of three native and one non-native species was significantly higher with burning, irrespective of whether or not the treatment had been mowed (Table 2). Native species were big bluestem (*Andropogon gerardii*) (1995; 62% cover), Indiangrass (*Sorghastrum nutans*) (1996; 25% cover), and leadplant (*Amorpha canescens*) (1995; 6% cover) (Fig. 2); the non-native species was red clover (*Trifolium pratense*) (1996; 5%). In contrast, four species were found which, for at least one year of the study, were significantly higher in areas from which burning had been excluded. These species were crown vetch (*Coronilla varia*) (1996; 3%), Scribner dichanthelium (1996; 6%), Kentucky bluegrass (1995; 21%), and red clover (1995; 13%). All but Scribner dichanthelium are non-native species.

Mowing.- No species were found that were significantly higher in mowed treatments. Overall, mowing alone resulted in significantly higher cover of smooth brome (*Bromus inermis* L.), an invasive, undesirable non-native grass (Fig. 3) irrespective of whether or not they had been burned. However, flowering spurge (*Euphorbia corollata*) (1996; 8%) and porcupine grass (1996; 2%) were two species in which maximum cover was significantly higher in the absence of mowing, although this treatment had also included burning. Canopy cover of finger coreopsis (*Coreopsis palmata*) (1995; 7%) and yellow sweet clover (*Melilotus officinalis*) (1996; 3%) were also significantly higher without mowing, but only in spring sampling periods.

Combined Burning and Mowing.- Blackseed plantain (*Plantago rugellii*) (1995; 6% cover and 1996; 1% cover) was the only species for which




Fig. 2. Mean canopy cover of big bluestem by treatment. Different letters at the top of each bar indicate significant differences ($P \leq 0.05$) between treatments based on Student-Newman-Keuls multicomparison test.




Fig. 2. Mean canopy cover of smooth brome by treatment. Different letters at the top of each bar indicate significant differences ($P \leq 0.05$) between treatments based on Student-Newman-Keuls multicomparison test.

maximum cover occurred with combined mowing and burning. However, in 1996, the year following burning, red clover also was significantly higher (8%) in the combined mowed-burned treatment areas. Both of these are non-native species.

DISCUSSION

In general, this study shows (1) that mowing and burning have the potential to affect substantially the floral composition of the tallgrass prairie, and (2) that burning is more suitable than mowing as a management practice for both maintaining native prairie species and for at least slowing the influx of aggressive, non-native species.

Many differences were noted in individual species and community-level responses to mowing and burning. Small sample size, experimental design that necessarily lacked close control of a substantial number of variables, and the unavoidable variability between individual treatments, such as specific timing and frequency, are but a few of the factors that complicate a reasonable interpretation of the results. Despite these factors, significant differences were noted that support the idea that different mowing and haying treatment of tallgrass prairie can affect species composition, results consistent with the literature (e.g. Dix 1960, McMurphy and Anderson 1965, Lovell *et al.* 1983).

One effect of treatment is the response to burning. Specifically, significant increases were noted for some of the dominant grasses and forbs, such as big bluestem, indiagrass, and leadplant (Table 2, Fig. 2) irrespective of mowing history. The effects on these grasses both support and contradict previous studies (Glenn-Lewin *et al.* 1990). In contrast, other native, but less common, grass species, in particular Scribner panicum, decreased in

response to burning. This opposite effect on native species indicated the difficulty in providing a single management for any site. It also supports the current trend towards varying management from year to year in order to facilitate ecosystem diversity (Solecki and Toney 1986, Fitzgerald and Tanner 1992).

The response of non-native species to burning is equally as important as is the response of native species (Ross and Vanderpoel 1991). Canopy cover of red clover, a non-native species, for example, decreased significantly immediately following burning but increased significantly the following year. Other species with lower cover in response to burning included Kentucky bluegrass, smooth brome, and crown vetch, the latter two both being particularly aggressive and undesirable, non-native species (Lovell *et al.* 1983). The recovery of red clover the year following burning, and the high proportion of smooth brome with mowing and in the absence of fire (Fig. 3), suggest that some form of regular, prescribed burning is necessary, if only to reduce the rate of invasion of these species, results consistent with studies by Evans (1983) and Fitzgerald and Tanner (1992).

Mowing, another management likely to be used in tallgrass prairie remnants, is viewed by some as a suitable alternative to fire (Ehrenreich 1959, Dale 1983, Mooberry 1984). This study suggests otherwise. In particular, none of the native species were found to be significantly higher in mowed areas, irrespective of burning history, whereas cover of smooth brome and blackseed plantain, both non-native species, was significantly higher with mowing. The response of blackseed plantain, however, occurred only when accompanied by burning. Not only did mowing not favor native species as a group, but it appears to have adversely affected them. For example, flowering

spurge and porcupine grass, both native species, were found to be significantly lower in mowed areas even when this treatment had also included burning. Canopy cover of finger coreopsis and yellow sweet clover, the latter a non-native ruderal species, was also significantly lower with mowing, but only in spring sampling periods.

Overall, my study suggests the importance of selecting management in maintaining the native tallgrass prairie plant diversity. Further, it supports the appropriate use of fire instead of mowing, both to favor native species and to limit the advance of non-native species. While many of the implications of my results are not new to the literature, they do provide yet more reinforcement to the argument favoring fire over mowing as a management tool where such application can be accommodated.

LITERATURE CITED

- Aldous, A.E. 1934. Effect of burning on Kansas bluestem pasture. Kansas Agriculture Experiment Station Bulletin 38. Kansas State University, Manhattan, Kansas.
- Anderson, K.L. 1965. Fire ecology - Some Kansas prairie forbs. *In* Proceedings of the Tall Timbers Fire Ecology Conference 4:153-160.
- Boettcher, J.F. and T.B. Bragg. 1989. Tallgrass prairie remnants of eastern Nebraska. Pages 1-7 *In* Proceedings of the Eleventh North American Prairie Conference, T. Bragg and J. Stubbendieck, editors. University of Nebraska, Lincoln, Nebraska.
- Bragg, T.B. 1982. Seasonal variations in fuel and fuel consumption by fires in a bluestem prairie. *Ecology* 63(1):7-11.
- Bragg, T.B. 1991. Implications for long-term prairie management from seasonal burning of loess hill and tallgrass prairies. Pages 20-24 *In* Proceedings of an international symposium. S.C. Nodvin and T.A. Waldrop, editors. Knoxville, TN.
- Bragg, T.B. 1995. Climate, soils and fire: The physical environment of North American grasslands. Pages 49-81 *In* The Changing Prairie, K. Keeler and A. Joern, editors. Oxford University Press, New York.
- Bragg, T.B. and L.C. Hulbert. 1976. Woody plant invasion of unburned Kansas bluestem prairie. *Journal of Range Management* 29:19-24.
- Conard, E.C. 1954. Effect of time of cutting on yield and botanical composition of prairie hay in southeastern Nebraska. *Journal of Range Management* 7(4):181-182.
- Curtis, J.T. and M.L. Partch. 1948. Effect of fire on competition between bluegrass and certain prairie plants. *The American Midland Naturalist* 39:437-443.
- Dale, E.E. 1983. The effects of mowing and burning on a restored prairie at Pea Ridge National Military Park, Benton County, Arkansas. Pp. 134-138 *In* Proceedings of the Eighth North American Prairie Conference, R. Brewer, editor. Western Michigan University, Kalamazoo, Michigan.
- Daubenmire, R.F. 1959. Canopy cover method of vegetation analysis. *Northwestern Scientist* 33:43-64.

- Dix, R.L. 1960. The effects of burning on the mulch structure and species composition of grasslands in western North Dakota. *Ecology* 41:49-56.
- Ehrenreich, J.H. 1959. Effect of burning and clipping on growth of native prairie in Iowa. *Journal of Range Management* 12(3):133-137.
- Ehrenreich, J.H. and J.M. Aikman. 1963. An ecological study of the effect of certain management practices on native prairie in Iowa. *Ecological Monographs* 33(2):113-130.
- Engle, D.M. 1988. Burning costs in Oklahoma rangelands. *Rangelands* 10(3):135-37.
- Evans, J.E. 1983. Literature review of management practices for smooth sumac (*Rhus glabra*), poison ivy (*Rhus radicans*) and other sumac species. *Natural Areas Journal* 3(1):16-26.
- Fitzgerald, S.M. and G.W. Tanner. 1992. Avian community response to fire and mechanical shrub control in south Florida. *Journal of Range Management* 45(4):396.
- Gehring, J.L. and T.B. Bragg. 1992. Woody vegetation of a disjunct bur oak (*Quercus macrocarpa* Michx.) forest in east-central Nebraska. *Transactions of the Nebraska Academy of Sciences* XX:41-46.
- Gibson, D.J., T.R. Seastedt, and J.M. Briggs. 1993. Management practices in tallgrass prairie: large- and small-scale experimental effects on species composition. *Journal of Applied Ecology* 247-255.
- Gillen, R.L. and R.W. McNew. 1987. Seasonal growth rates of tallgrass prairie after clipping. *Journal of Range Management* 40:342-345.
- Glenn-Lewin, D.C., L.A. Johnson, T.W. Jurik, A. Akey, M. Leoschke, and T. Rosburg. 1990. Fire in central North American grasslands: Vegetative reproduction, seed germination, and seedling establishment. Pages 28-45 in *Fire in North American tallgrass prairies* (S.L. Collins and L.L. Wallace, editors). University of Oklahoma Press, Norman, OK.
- Great Plains Flora Association. 1986. *Flora of the Great Plains*. University Press of Kansas, Lawrence, KS.

- Hadley, E.B. and B.J. Kieckhefer. 1963. Productivity of two prairie grasses in relation to fire frequency. *Ecology* 44:389-395.
- Hadley, E.B. 1970. Net productivity and burning response of native eastern North Dakota prairie communities. *The American Midland Naturalist* 84:121-135.
- Hanes, C.R. and F.N. Hanes. 1947. *Flora of Kalamazoo County, Michigan*. Schoolcraft, Michigan.
- Hayden, A. and J.M. Aikman. 1949. Considerations involved in the management of prairie reserves. *Iowa Academy of Science* 56:133-142.
- Hensel, R.L. 1923. Recent studies on the effect of burning on grassland vegetation. *Ecology* 4:183-188.
- Hover, E.I. and T.B. Bragg. 1981. Effect of season of burning and mowing on an eastern Nebraska *Stipa-Andropogon* prairie. *The American Midland Naturalist* 105:13-18.
- Hulbert, L.C. 1969. Fire and litter effects in undisturbed bluestem prairie in Kansas. *Ecology* 50(5):874-877.
- Hulbert, L. 1986. Fire effects on tallgrass prairie. Pages 138-142. *In* Proceedings of the Ninth North American Prairie Conference, G.K. Clamby and R.H. Pemble, editors. Tri-College University Center for Environmental Studies, Fargo, North Dakota.
- Hulbert, L. 1988. Causes of fire in the tallgrass prairie. *Ecology* 69(1):46-58.
- Kerr, K. and J. White. 1981. A volunteer-supported effort to find an preserve prairie and savanna remnants in Illinois cemeteries. Page 6. *In* Proceedings of the Sixth North American Prairie Conference, R.L. Stuckey and K.J. Reese, eds. Ohio State University, Columbus, Ohio.
- Kline, V. 1986. Response to sweet clover (*Melilotus alba* Desr.) and associated prairie vegetation to seven experimental burning and mowing treatments. Pages 149-152 *In* The Prairie: past, present and future. Proceedings of the Ninth North American Prairie Conference, G.K. Clamby and R.H. Pemble, editors. Tri-College University Center for Environmental Studies, Fargo, North Dakota.

- Klopatek, J.M., R.J. Olson, C.J. Emerson, and J.L. Jones. 1979. Land-use conflicts with natural vegetation in the United States. *Environmental Conservation* 6:191-199.
- Knapp, A.K. and T.R. Seastedt. 1986. Detritus accumulation limits productivity of tallgrass prairie. *BioScience* 36(10):662-668.
- Kucera, C.L. and M. Koelling. 1964. The influence of fire on composition of central Missouri prairie. *American Midland Naturalist*. 72:142-147.
- Küchler, A. W. 1964. Potential natural vegetation of the conterminous United States. *American Geographical Soc. Special Publ.* 36.
- Küchler, A.W. 1985. Potential national vegetation. *National Atlas of the United States of America, Department of the Interior, U.S. Geological Survey. Reston, Virginia. Map.*
- Launchbaugh, J.L. and E.E. Owensby. 1978. Kansas Rangelands: Their management based on a half century of research. *Kansas Agriculture Experiment Station Bulletin 622, Kansas State University, Manhattan, Kansas.*
- Leach, M.K. and T.J. Givnish. 1996. Ecological determinants of species loss in remnant prairies. *Science* 273(5281):1555-1558.
- Leenhouts, B. 1997. Presettlement fire and emission production estimates: A framework for understanding potential system change. Pages 236-241 *In Environmental Regulation & Prescribed Fire: Legal and Social Challenges*, D.C. Bryan, editor. Center for Professional Development, Florida State University.
- Lovell, D.L., R.A. Henderson, and E.A. Howell. 1983. The response of forb species to seasonal timing of prescribed burns in remnant Wisconsin prairies. Pages 11-15 *In Proceedings of the Eighth North American Prairie Conference*, R. Brewer, editor. Western Michigan University, Kalamazoo, Michigan.
- McMurphy, W.E. and K.L. Anderson. 1965. Burning Flint Hills range. *Journal of Range Management* 18:265-268.
- Mooberry, F.M. 1984. Meadow response to April, June, July mowing (Pennsylvania). *Restoration and Management Notes* 2:27-28.

- Morris, M.G. 1979. Responses of grassland invertebrates to management by cutting: Heteroptera. *Journal of Applied Ecology* 16:417-432.
- National Oceanic and Atmospheric Administration. 1989. Local climatological data: Annual summary with comparative data - Omaha (North), Nebraska. National Climatic Data Center, Asheville, NC.
- Old, S.M. 1969. Microclimate, fire, and plant production in an Illinois prairie. *Ecological Monographs* 39:355-384.
- Pemble, R.H., G.L. Van Amburg, and L. Mattson. 1981. Intraspecific variation in flowering activity following a spring burn on a northwestern Minnesota prairie. Pages 235-240 *In* The Prairie Peninsula-In the "shadow" of Transeau: Proceedings of the Sixth North American Prairie Conference, R.L. Stuckey and K.J. Reese, editors. Ohio State University, Columbus, Ohio.
- Pyne, Stephen J. 1982. Fire Primeval. *The Sciences* 22(6):14-20.
- Robocker, C.W. and B.J. Miller. 1955. Effects of clipping, burning, and competition on establishment and survival of some native grasses in Wisconsin. *Journal of Range Management* 8:117-121.
- Ross, L.M. and T. Vanderpoel. 1991. Mowing encourages establishment of prairie species (Illinois). (Abstract). *Restoration and Management Notes* 9(1):34-35.
- Samson, F. and F. Knopf. 1994. Prairie conservation in North America. *Bioscience* 44(6):418-21.
- SAS Institute Inc. 1985. SAS user's guide: Statistics, version 5 edition. Cary, North Carolina: SAS Institute Inc.
- Schwegman, J.E. and W.E. McClain. 1985. Vegetative effects and management implications of a fall prescribed burn in an Illinois hill prairie. *Natural Areas Journal* 5(3):4-8.
- Solecki, M.K., J.B. Taft, L.A. Cook, and P.S. Haverland. 1986. Vegetational composition of three Missouri tallgrass prairies with reference to past management. Missouri Department of Conservation, Jefferson City, Missouri.

- Solecki, M.K. and T. Toney. 1986. Characteristics and management of Missouri's public prairies. Pages 168-171 *In* Proceedings of the Ninth North American Prairie Conference, G.K. Clambey and R.H. Pemble, editors. Tri-College University Center for Environmental Studies, Fargo, North Dakota.
- Tomanek, G.W. 1948. Pasture types of western Kansas in relation to the intensity of utilization in past years. *Transcripts of the Kansas Academy of Science* 51:171-196.
- Towne, G. and C. Owensby. 1984. Long-term effects of annual burning at different dates in ungraded Kansas tallgrass prairie. *Journal of Range Management* 37(5):392-397.
- Weaver, J.E. and T.J. Fitzpatrick. 1934. The prairie. *Ecological Monographs* 4(1):109-295.
- Willson, G.D. and J. Stubbendieck. 1996. Suppression of smooth brome by atrazine, mowing and fire. *The Prairie Naturalist* 28(1):13-20.
- Zar, J.H. 1984. *Biostatistical Analysis*. Prentice-Hall, Inc. Englewood Cliffs, NJ.

Appendix

Appendix Table 1. Shannon Weiner Diversity Index (Matrix) Values of all species, at eight Iowa cemetery sites. SD=significant difference between sites, NSD=no significant difference between sites. Sites have the following treatments: 1,2,3 are burned, 4,5,6 are mowed and burned, 7,8 are mowed.

	1	2	3	4	5	6	7	8
1		NSD t=1.460 t _{0.05(2)} 371 =1.967	SD t=3.853 t _{0.05(2)} 713 =1.963	SD t=2.864 t _{0.05(2)} 349 =1.967	SD t=3.022 t _{0.05(2)} 342 =1.967	SD t=5.241 t _{0.05(2)} 362 =1.967	NSD t=1.482 t _{0.05(2)} 104 =1.983	NSD t=5.828 t _{0.05(2)} 357 =1.967
2			SD t=6.004 t _{0.05(2)} 12 =2.179	NSD t=1.603 t _{0.05(2)} 352 =1.967	SD t=5.166 t _{0.05(2)} 413 =1.966	SD t=4.201 t _{0.05(2)} 368 =1.966	NSD t=3.444 t _{0.05(2)} 86 =1.988	NSD t=8.822 t _{0.05(2)} 360 =1.967
3				SD t=7.351 t _{0.05(2)} 344 =1.967	NSD t=1.144 t _{0.05(2)} 511 =1.965	SD t=10.069 t _{0.05(2)} 366 =1.967	SD t=4.821 t _{0.05(2)} 76 =1.992	SD t=4.699 t _{0.05(2)} 357 =1.967
4					SD t=6.595 t _{0.05(2)} 334 =1.967	SD t=2.484 t _{0.05(2)} 347 =1.967	NSD t=9.172 t _{0.05(2)} 95 =1.985	SD t=2.364 t _{0.05(2)} 341 =1.967
5						SD t=9.403 t _{0.05(2)} 265 =1.969	SD t=4.141 t _{0.05(2)} 71 =1.993	SD t=3.861 t _{0.05(2)} 347 =1.967
6							SD t=3.587 t _{0.05(2)} 97 =1.984	SD t=4.830 t _{0.05(2)} 364 =1.967
7								NSD t=1.022 t _{0.05(2)} 96 =1.985