

Student Work

---

7-1-1983

**Vegetational changes under isolated *Juniperus virginiana* in an eastern Nebraska bluestem prairie.**

Janet L. Gehring

Follow this and additional works at: <https://digitalcommons.unomaha.edu/studentwork>

---

**Recommended Citation**

Gehring, Janet L., "Vegetational changes under isolated *Juniperus virginiana* in an eastern Nebraska bluestem prairie." (1983). *Student Work*. 3384.

<https://digitalcommons.unomaha.edu/studentwork/3384>

This Thesis is brought to you for free and open access by DigitalCommons@UNO. It has been accepted for inclusion in Student Work by an authorized administrator of DigitalCommons@UNO. For more information, please contact [unodigitalcommons@unomaha.edu](mailto:unodigitalcommons@unomaha.edu).



VEGETATIONAL CHANGES UNDER ISOLATED  
JUNIPERUS VIRGINIANA IN AN EASTERN  
NEBRASKA BLUESTEM PRAIRIE

A Thesis

Presented to the

Department of Biology

and the

Faculty of the Graduate College

University of Nebraska

In Partial Fulfillment

of the Requirements for the Degree

Master of Arts

University of Nebraska at Omaha

by

Janet L. Gehring  
July 1983

UMI Number: EP74986

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI EP74986

Published by ProQuest LLC (2015). Copyright in the Dissertation held by the Author.

Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code



ProQuest LLC.  
789 East Eisenhower Parkway  
P.O. Box 1346  
Ann Arbor, MI 48106 - 1346

## THESIS ACCEPTANCE

Accepted for the faculty of the Graduate College,  
University of Nebraska, in partial fulfillment of the  
requirements for the degree Master of Arts, University of  
Nebraska at Omaha.

## Committee

Name	Department
<i>David M. Sutherland</i>	<i>Biology</i>
<i>Larry J. Stephens</i>	<i>Math/CS</i>

*Thomas B. Bragg*  
Chairman

*11 July 1983*  
Date

## ACKNOWLEDGMENTS

I would like to thank the members of my graduate committee for their valuable input and suggestions throughout this project: Dr. Thomas Bragg for suggesting the project, assisting with the manuscript, and for continuous advice, understanding, and patience; Dr. David Sutherland for plant identifications, assisting with manuscript figures and manuscript review; Dr. Larry Stephens for statistical advice and manuscript review. I would also like to thank Dr. William DeGraw for help with seminar slides.

I would like to express my appreciation to my fellow graduate students, professors, co-workers, and friends for encouragement and support, not only during this project, but also during the entire course of my graduate studies. I am especially grateful to Betty Carlson for typing this manuscript.

Most of all, my heartfelt gratitude is extended to my family for their emotional and financial support. They have always been there to provide a solid foundation for my life and all my endeavors.

## TABLE OF CONTENTS

	PAGE
THESIS ACCEPTANCE . . . . .	ii
ACKNOWLEDGEMENTS . . . . .	iii
LIST OF TABLES . . . . .	v
LIST OF FIGURES . . . . .	vi
ABSTRACT . . . . .	vii
INTRODUCTION . . . . .	1
METHODS AND MATERIALS . . . . .	3
STUDY SITE . . . . .	3
DATA COLLECTION . . . . .	3
DATA ANALYSIS . . . . .	3
RESULTS . . . . .	7
DISCUSSION . . . . .	12
LITERATURE CITED . . . . .	15
APPENDIX . . . . .	18

## LIST OF TABLES

TABLE		PAGE
1.	Mean percent canopy cover + SE for species with significant differences in distribution between successional plots . . . . .	8
Appendix Table I.	Floristics of study plots . . . . .	19
Appendix Table II.	Between-site differences separated by plot and aspect . . . . .	25
Appendix Table III.	Analysis of variance (ANOVA) on transformed-percent-cover . . . . .	26

LIST OF FIGURES

FIGURE	PAGE
1. Mean percent canopy cover for <u>Andropogon scoparius</u> and <u>Poa pratensis</u> . . . . .	9
Appendix Figure 1. Regression of transformed data of combined other grasses onto canopy radius . . . . .	27
Appendix Figure 2. Regression of transformed data of <u>Poa pratensis</u> onto canopy radius . . . . .	28

6



## ABSTRACT

Changes in species composition of a bluestem prairie were evaluated under isolated, 10-22 year old Juniperus virginiana (Eastern Red Cedar) in eastern Nebraska. Percent canopy cover was measured in full-canopy, canopy-edge, and open-canopy plots which were located along transects oriented to the cardinal compass directions. A total of 53 species were recorded. Andropogon scoparius and A. gerardii dominated open-canopy and canopy-edge plots; Poa pratensis and A. scoparius dominated full-canopy plots. Canopy cover of most prairie species decreased under full canopy but Poa pratensis and Carex spp. increased. There was a positive correlation between tree size and both the decrease of prairie species and the increase of Poa pratensis under a full-canopy. Similarly, aspect was significantly correlated to changes in species composition under a full-canopy. For those species with significant differences between cardinal directions, canopy cover was lowest on the north and east transects. The results of this study show that invasion of woody plants into native prairie rapidly affects the composition of native grassland species.

## INTRODUCTION

Fire was a major environmental factor affecting the bluestem prairie (Küchler 1964) that once dominated much of the eastern portion of the North American grasslands. Extensive reviews of the effects of fire have been conducted by Daubenmire (1968) and Vogl (1974). Among the many effects noted is that fire inhibits woody plant invasion of prairie by directly killing or injuring many species of trees and shrubs (Kucera 1960, Blan 1970, Bragg and Hulbert 1976) and indirectly by reducing available soil moisture for seedlings (Hulbert 1969).

Grassland species can be divided into two main groups according to their mode of carbon fixation during photosynthesis:  $C_3$ , or cool-season and  $C_4$ , or warm-season (Williams 1974, Waller and Lewis 1979).  $C_4$  species are thought to have a competitive advantage under conditions of high light, high temperature, and decreased moisture (Bjorkman 1971, Black 1971). Although  $C_3$  and  $C_4$  species often grow in the same habitat, most of the dominant species of bluestem prairies in eastern Nebraska possess the  $C_4$  pathway (Waller and Lewis 1979). Moderation of the microenvironment by invading woody plants may cause the  $C_4$  species to lose their advantage due to the effects of shading. Thus, under these modified conditions, species composition should reflect an increasing proportion of  $C_3$

plants under the tree canopy. In addition, one of the major non-native invaders of bluestem prairie, Poa pratensis L. (Kentucky bluegrass) (Weaver 1965), is a C<sub>3</sub> species which may also be favored by the conditions created by invading woody plants. Changes in plant species composition under invading woody plants has been studied in other grassland ecosystems (Arnold 1964, Jameson 1966) as well as under trees in savanna ecosystems (Bray 1955, Parker and Muller 1982). No such studies, however, have been documented for the bluestem prairie of eastern Nebraska.

The objective of this study was (1) to document changes in species composition under invading trees and (2) to test the hypothesis that there is a correlation between the invasion of woody species and changes in species composition that reflect the effects of different photosynthetic pathways.

## METHODS AND MATERIALS

## Study Site:

The study was conducted on a native prairie located on bluffs adjacent to the Platte River Valley in southeastern Sarpy County, Nebraska (W1/2Sec34 T13N R10E). The bluffs are generally west-facing with 17 to 30 percent slopes. The loessal soils are Ida silt loams, low in organic matter and nitrogen, mildly to moderately alkaline, and well-drained (Bartlett 1975). Precipitation averages 71 cm annually with 78% of the precipitation falling during the growing season. Average temperatures range from -6 C in January to 24 C in July (U.S. Department of Commerce 1981).

The study area was chosen because (1) the prairie was mostly undisturbed and not presently or recently grazed, and (2) isolated Juniperus virginiana L. (Eastern Red Cedar), a common early invader of tallgrass prairie, was found on the slopes and ridgetops.

## Data Collection:

Six separate south- or southwest-facing slopes were selected as study sites, based on their having representative native prairie vegetation and an appropriate number and size of isolated trees. Trees were considered isolated if neighboring trees did not cast shadows on them during the majority of the day. Within each study area, six trees of approximately the same size were selected and marked with

surveyor's flagging. An increment core was extracted near the base of each tree to provide an estimate of the time of its establishment and hence a measure of the length of time that microclimatic changes had affected plant composition. Ring counts were made on each core using standard techniques (Stokes and Smiley 1968). Other data recorded at each tree included tree canopy radius, the distance from the main tree stem to canopy edge, and percent slope.

Four transects were established at each tree, one for each of the cardinal compass headings. Cardinal direction was considered to be an important factor, not only because of solar angle, but also because of steep slopes at the study area. Three 20 x 50 cm plots were located along each transect with the long axis parallel to the canopy edge. One plot was located 70 cm from the main tree stem, one at the edge of the canopy, and one 70 cm outside of the canopy. These represent full-canopy, canopy-edge, and open-canopy plots respectively and, in combination, are termed successional plots. Open-canopy plots were considered to be the control. Each plot was evaluated once between July 24 and August 6, 1982. Data recorded for each plot included total vegetative cover, total grass cover, total forb cover, and total woody plant cover, as well as percent canopy cover by species. Canopy coverage categories were 0-5%, 5-25%, 50-75%, and 95-100% (Daubenmire 1959). Data were analyzed using mid-point values for each coverage category. Dominant

species are defined as (1) the two species with the highest mean canopy cover for each group of successional plots and (2) any additional species within five percent canopy cover of these species.

Plant identifications were verified at the University of Nebraska at Omaha Herbarium (OMA). Common and scientific names are from Great Plains Flora Association (1977), and Sutherland (In press).

#### Data Analysis:

Vegetative diversity for each group of successional plots was calculated using the Shannon-Wiener Diversity Function ( $H'$ ) where  $H' = -\sum p_i \log p_i$  and  $p_i$  = the relative canopy coverage value of each species; high  $H'$  values indicate high diversity (Krebs 1978). Pearson's Correlation Coefficient,  $r$ , was used to determine the strength of linear relationships between two variables (Ott 1977). The Kruskal-Wallis test, a non-parametric procedure, was used for statistical analysis on canopy cover data of individual species because of unequal variance in the data (Ott 1977).

Transformation of the data was used to reduce variability and normalize canopy cover data. This involved dividing total grass cover by mean canopy cover of Poa pratensis and by the sum of mean values for all other grasses. Data from this manipulation are termed "transformed-percent-cover." Grasses that were combined included Andropogon gerardii Vitman (Big Bluestem), A.

scoparius Michx. (Little Bluestem), Bouteloua curtipendula (Michx.) Torr. (Side-oats Grama), B. hirstua Lag. (Hairy Grama), Dichanthelium oligosanthes var. scribnerianum (Nash.) Gould (Scribner Panicum), D. wilcoxianum (Vasey) Freckman (Wilcox Panicum), Festuca octoflora Walt. (Six-weeks Fescue), Koeleria pyramidata (Lam.) Beauv. (Junegrass), Muhlenbergia cuspidata (Torr. in Hook) Rydb. (Plains Muhly), Paspalum setaceum Michx., and Sorghastrum nutans (L.) Nash (Indian grass). Poa pratensis was analyzed separately because it was the only grass that increased significantly under the tree canopy. Transformed-percent-cover of Poa pratensis and of other combined grasses were tested using an Analysis of Variance (ANOVA) (Nie et al., 1975). All statistical hypotheses were considered significant if  $p \leq 0.05$ .

## RESULTS

A total of 53 species were recorded in 432 plots (Appendix Table I). Fourteen species averaged greater than 10% and 25 species averaged less than 1% frequency.

Andropogon scoparius and A. gerardii dominated in open-canopy and canopy-edge plots; Poa pratensis and Andropogon scoparius dominated in full-canopy plots (Table 1). In addition, eight species of woody plant seedlings were recorded in the study plots: Acer negundo L. (Box Elder), Cornus drummondii Mey. (Rough-leaved Dogwood), Fraxinus pennsylvanica Marsh. (Green Ash), Juniperus virginiana, Quercus macrocarpa Michx. (Bur Oak), Ribes missouriense Nutt. (Missouri Gooseberry), Symphoricarpos orbiculatus Moench (Buckbrush), and Ulmus spp. (Elm).

Canopy cover of 16 species differed significantly between full-canopy, canopy-edge, and open-canopy plots. For 11 of these species, the lowest mean percent cover occurred in full-canopy plots; values averaged highest in full-canopy plots for Poa pratensis, Carex spp., and Juniperus virginiana (Table 1). Changes in Andropogon scoparius, a warm-season, C<sub>4</sub> grass, and Poa pratensis, a cool-season, C<sub>3</sub> grass, represent the major patterns observed (Fig. 1). Dichanthelium oligosanthos var. scribnerianum, also a C<sub>3</sub> species like Poa pratensis, decreased under the tree canopy. A regression of transformed-percent-cover of the full-canopy



Table 1. Mean percent canopy cover + SE for species with significant differences ( $p < 0.05$ ) in distribution between successional plots based on Kruskal-Wallis Analysis of Variance:  $tr < 0.5\%$  cover.  $H' =$  Shannon-Wiener function for combined successional plots. Values represent combined data for all six study sites.

FLORISTICS	SUCCESSIONAL PLOTS		
	Full-Canopy	Canopy-Edge	Open-Canopy
SPECIES DIVERSITY ( $H'$ )	2.586	2.706	2.622
SPECIES:			
<u>Andropogon scoparius</u> (Little Bluestem)	10.3+1.42	38.4+2.31	43.7+2.34
<u>Poa pratensis</u> (Kentucky Bluegrass)	19.1+1.54	10.1+1.18	6.4+0.91
<u>Aster ericoides</u> (White Aster)	tr	3.6+0.49	5.0+0.62
<u>Bouteloua curtipendula</u> (Side-oats Grama)	3.9+0.84	8.6+1.24	5.6+0.91
<u>Dichanthelium oligosanthos</u> var. <u>scribnerianum</u> (Scribner Panicum)	0.8+0.16	2.8+0.37	2.6+0.48
<u>Andropogon gerardii</u> (Big Bluestem)	3.6+0.84	15.4+2.16	13.8+2.03
<u>Ambrosia psilostachya</u> (Western Ragweed)	tr	0.8+0.16	0.8+0.16
<u>Carex</u> spp. (Sedge)	1.2+0.27	tr	tr
<u>Solidago</u> spp. (Goldenrod)	tr	0.6+0.05	0.6+0.07
<u>Bouteloua hirsuta</u> (Hairy Grama)	tr	2.3+0.67	2.3+0.20
<u>Linum rigidum</u> (Flax)	tr	0.4+0.07	0.6+0.12
<u>Lygodesmia juncea</u> (Skeleton Weed)	tr	tr	0.8+0.19
<u>Petalostemum purpureum</u> (Purple Prairie Clover)	tr	0.6+0.19	1.1+0.30
<u>Juniperus virginiana</u> seedlings (Eastern Red Cedar)	0.8+0.21	tr	tr
<u>Calylophus serrulatus</u> (Yellow Evening Primrose)	tr	tr	tr
<u>Hedeoma hispida</u> (Rough Penneyroyale)	tr	tr	tr




Figure 1. Mean percent canopy cover for Andropogon scoparius and Poa pratensis. Horizontal lines represent mean values, vertical lines connect maximum and minimum values, and vertical bars represent one standard error.

plots onto canopy radius, which reflects the length of time that full-canopy plots have been affected, further documents the sequential loss of prairie grasses ( $r=-.597$ ,  $p=.005$ ) and the increase of *Poa pratensis* ( $r=.555$ ,  $p=.005$ ) (Appendix Fig. 1-2). No correlation was observed for open-canopy and canopy-edge plots.

Significant differences were found between sites for all successional plot groups indicating that plant species composition varied somewhat between study sites despite their topographic similarity.

The effect of plot direction from the main tree stem was analyzed in several different ways. Firstly, of the six most abundant species, the greatest number of significant, between-site variations was found for west aspect successional plots (Appendix Table II). Secondly, between-plot differences were separated by direction to assess the contribution of direction to between-plot differences for each species. West transects had the least number of significant differences (5), with increasing numbers of differences for south (8), north (11), and east (12). Thirdly, without transformation of data, Kruskal-Wallis analysis of variance resulted in relatively few significant between-aspect differences for each of the three plot groups for individual species. For those species, canopy cover was lowest on the north and east transects. Fourthly, after data transformation, aspect was found to be

a significant factor for both Poa pratensis ( $p=0.02$ ) and for other combined grasses ( $p=0.005$ ). Interaction between plot and aspect was also significant for both groups (Appendix Table III).

The approximate age of trees chosen for the study ranged from 10-22 yr. Canopy radius increased with tree age ( $r=0.523$ ,  $p=.005$ ).

## DISCUSSION

Grassland species are substantially affected by invading trees with the main effect occurring after complete coverage by the tree canopy. Two distinct patterns were found, one negative and one positive. All native prairie species decreased under full canopy with the exception of Carex spp. which increased. Poa pratensis, a non-native C<sub>3</sub> species, also increased as the tree canopy increased. This response has been reported to occur under other conditions such as drought and overgrazing where native prairie species decrease in abundance (Weaver 1965) thus supporting the idea that its increase may be a response to release from competition.

Tree size is an important factor particularly as it relates to time period of canopy influence. There is a sequential loss of native prairie species as they are covered by the canopy for longer periods of time. The largest trees had low vegetative cover under full-canopy, indicating that eventually no grassland species can grow under the environmental conditions that exist directly under the tree. As the tree canopy increases its influence, the area in which native prairie species are eliminated will increase.

Photosynthetic pathway was not an important predictive factor for negative or positive response under full-canopy. Of the two C<sub>3</sub> species that had sufficiently large

populations to allow for statistical comparison, Poa pratensis increased under full-canopy whereas Dichanthelium oligosanthos var. scribnerianum decreased. One other native C<sub>3</sub> grass species, Koeleria pyramidata (Junegrass), was recorded in low abundance in canopy-edge and open-canopy plots and was absent under full-canopy. Other ecological factors may account for these different responses, of which some possible explanations include phytotoxicity of Juniperus virginiana (Rice 1974), effects of tree litter (Jameson 1963, 1966, Rice 1974), and changes in water availability due to rainfall interception (Anderson et al. 1969). Any or all of these factors may act in conjunction with shading effects to cause the observed response to canopy closure. At least for some species these factors may mask the contribution of the type of photosynthetic pathway to declines in species cover.

The effects of aspect, or the cardinal direction from the main tree stem, were observed for a limited number of species due to high variation in the data. For those species with significant differences, canopy cover was lower on the north and east transects when compared to south and west transects, indicating solar angle was a factor. The combined data analysis of variance reinforced the importance of aspect.

The general results of this study show that, even when woody plant invasion is restricted to isolated trees, there is an effect on species composition of the prairie that is

affected by aspect and increases as invasion continues. Further, the presence of woody plant seedlings in the study plots indicates that a single invader, such as Juniperus virginiana, provides a favorable environment for the establishment of other woody plants. In combination, the increase in woody plants and the decline in prairie species reaches a point where the one-time prairie will be irretrievably changed. Ecosystem managers need to be aware of these effects and the rate of changes in species composition so as to prevent excessive invasion and coincident loss of native species from areas intended to be maintained in their pristine state.

## LITERATURE CITED

- Anderson, R. C., O. L. Loucks, and A. M. Swain. 1969. Herbaceous response to canopy cover, light intensity, and throughfall precipitation in coniferous forests. *Ecology* 50: 255-263.
- Arnold, J. F. 1964. Zonation of understory vegetation around a juniper tree. *Journal of Range Management* 17: 41-42.
- Bartlett, P. A. 1975. Soil Survey of Douglas and Sarpy Counties, Nebraska. U.S. Department of Agriculture S. C. S. 79 pp.
- Bjorkman, O. 1971. Comparative photosynthetic CO<sub>2</sub> exchange in higher plants. Pages 18-22 in M. D. Hatch, C. B. Osmond, and R. O. Slayter. *Photosynthesis and photorespiration*. Wiley-Interscience, New York.
- Black, C. 1971. Ecological implications of dividing plants into groups with distinct photosynthetic production capacities. *Advances in Ecological Research* 7: 87-114.
- Blan, K. 1970. Evaluation of eastern red cedar (Juniperus virginiana L.) infestations in the northern Kansas Flint Hills. M. S. Thesis. Kansas State University, Manhattan. 29 pp.
- Bragg, T. B. and L. C. Hulbert. 1976. Woody plant invasion of unburned Kansas bluestem prairie. *Journal of Range Management* 29: 19-24.
- Bray, J. R. 1955. The savanna vegetation of Wisconsin and application of the concepts order and complexity to the field of ecology. Unpublished Ph.D. Dissertation. University of Wisconsin, Madison. 251 pp.
- Daubenmire, R. 1959. A canopy-coverage method of vegetational analysis. *Northwest Science* 33: 43-64.
- Daubenmire, R. 1968. Ecology of fire in grasslands. *Advances in Ecological Research* 5: 209-266.
- Great Plains Flora Association. 1977. Atlas of the flora of the Great Plains. Iowa State University Press, Ames. 600 pp.



- Hulbert, L. C. 1969. Fire and litter effects in undisturbed bluestem prairie in Kansas. *Ecology* 50: 874-877.
- Jameson, D. A. 1963. Plant extracts retard growth of grass seedlings. In *Rocky Mountain Forest and Range Experimental Station, ed. Annual Report for 1963.* USDA, Fort Collins, Colorado.
- Jameson, D. 1966. Pinyon-juniper litter reduces growth of blue grama. *Journal of Range Management* 19: 214-217.
- Krebs, C. 1978. *Ecology: the experimental analyses of distribution and abundance*, 2nd Edition. Harper and Row, New York. 694 pp.
- Kucera, C. L. 1960. Forest encroachment in native prairie. *Iowa State Journal of Science* 34: 635-639.
- Küchler, A. W. 1964. Potential natural vegetation of the conterminous United States. *American Geographical Society Special Publication No. 36.* 116 pp.
- Nie, N., C. H. Hull, J. G. Jenkins, K. Steinbrenner, and D. H. Bent. 1975. *Statistical package for the social sciences (SPSS) manual*, 2nd Edition. McGraw-Hill Book Co., Belmont, California. 724 pp.
- Ott, L. 1977. *An introduction to statistical methods and data analysis.* Duxbury Press: North Scituate, Massachusetts. 730 pp.
- Parker, V. T. and C. H. Muller. 1982. Vegetation and environmental changes beneath isolated live oak trees (*Quercus agrifolia*) in a California annual grassland. *American Midland Naturalist* 107: 69-81.
- Rice, E. L. 1974. Roles of allelopathy in patterning of vegetation and creation of bare areas. Pages 126-173 in *E. L. Rice. Allelopathy.* Academic Press, New York.
- Stokes, M. A. and T. L. Smiley. 1968. *An introduction to tree-ring dating.* The University of Chicago Press, Chicago. 73 pp.
- Sutherland, D. In press. In Great Plains Flora Association. *Flora of the Great Plains.* Iowa State University Press, Ames.

- U.S. Department of Commerce. 1981. Local climatological data monthly summary, Omaha, Nebraska. National Oceanographic and Atmospheric Administration. Environmental Data, Sve., Asheville, North Carolina.
- Vogl, R. J. 1974. Effects of fire on grasslands, pp. 139-194. In T. T. Kozlowski and C. E. Ahlgren, eds. Fire and ecosystems. Academic Press, New York.
- Waller, S. S. and J. K. Lewis. 1979. Occurrence of C<sub>3</sub> and C<sub>4</sub> photosynthetic pathways in North American grasses. Journal of Range Management 32: 12-28.
- Williams, G. J. 1974. Photosynthetic adaptation to temperature in C<sub>3</sub> and C<sub>4</sub> grasses. Plant Physiology 54: 709-711.
- Weaver, J. E. 1965. Native vegetation of Nebraska. University of Nebraska Press, Lincoln. 185 pp.

APPENDIX

Appendix Table I. Floristics of study plots, 1 = full canopy, 2 = canopy edge, 3 = open canopy, N = north, S = south, E = east, W = west, TR < 0.1% for relative frequency.

Species:	% Rel. Freq.	Mean percent canopy cover $\pm$ SE											
		1-N	2-N	3-N	1-S	2-S	3-S	1-E	2-E	3-E	1-W	2-W	3-W
<u>Andropogon scoparius</u>	82	5.1 +1.63	39.8 +5.16	45.2 +4.65	13.9 +3.26	35.6 +4.39	45.8 +4.86	7.0 +2.45	40.9 +4.58	45.5 +4.87	15.3 +3.42	37.4 +4.42	38.5 +4.37
<u>Poa pratensis</u>	74	19.2 +2.90	11.6 +2.11	6.58 +1.64	18.7 +3.06	6.5 +2.25	3.9 +1.29	19.7 +2.99	9.4 +2.01	7.56 +2.20	18.9 +3.47	12.9 +2.93	7.5 +2.02
<u>Aster ericoides</u>	56	.3 + .12	2.9 + .74	4.3 +1.31	.8 + .17	4.6 +1.02	4.6 + .95	.3 + .12	3.7 +1.24	6.0 +1.40	.6 + .16	3.1 + .82	5.0 +1.32
<u>Bouteloua curtipendula</u>	55	1.7 + .69	2.3 + .78	5.6 +1.29	7.4 +2.75	14.3 +3.47	6.2 +2.05	1.4 + .58	8.5 +2.04	4.6 +1.57	5.1 +1.58	9.5 +2.52	7.8 +2.25
<u>Dichanthelium oligosanthos</u> var. <u>scribnerianum</u>	51	.5 + .15	2.1 + .55	2.7 + .95	.6 + .15	3.4 + .89	2.1 + .68	1.1 + .43	2.9 + .74	3.8 +1.23	1.1 + .43	2.6 + .76	1.78 + .57
<u>Andropogon gerardii</u>	33	3.6 +1.18	19.7 +4.49	13.2 +3.43	4.7 +2.25	9.5 +3.34	8.4 +2.9	2.2 + .87	17.2 +4.62	16.5 +4.56	5.2 +2.04	15.2 +4.73	16.9 +5.00
<u>Ambrosia psilostachya</u>	26	.2 + .09	.8 + .13	.8 + .15	.2 + .11	1.0 + .43	.8 + .43	.3 + .12	1.2 + .43	1.1 + .43	.3 + .13	.7 + .16	.7 + .16
<u>Carex</u> spp.	23	2.1 + .78	1.0 + .43	.5 + .12	1.3 + .58	.2 + .11	.3 + .13	.9 + .43	.3 + .12	.4 + .14	.6 + .16	.6 + .15	1.2 + .58
<u>Solidago</u> spp.	20	.1 + .08	.28 + .12	.6 + .16	.4 + .13	2.8 + .43	.5 + .15	0.0 + .00	.9 + .43	.6 + .15	.2 + .11	.6 + .15	.5 + .15

Appendix Table I. Floristics of study plots (continued). Mean percent canopy cover  $\pm$  SE, and relative frequency.

Species:	% Rel. Freq.	Mean percent canopy cover $\pm$ SE											
		1-N	2-N	3-N	1-S	2-S	3-S	1-E	2-E	3-E	1-W	2-W	3-W
<u>Bouteloua hirsuta</u>	17	1.0 $\pm$ 1.06	0.4 $\pm$ .42	2.3 $\pm$ .87	0.4 $\pm$ .42	7.0 $\pm$ 2.41	4.1 $\pm$ 1.33	0.5 $\pm$ .42	0.5 $\pm$ .42	0.6 $\pm$ .42	0.1 $\pm$ .06	1.2 $\pm$ .58	2.4 $\pm$ .86
<u>Linum rigidum</u>	16	0.0 $\pm$ .00	0.1 $\pm$ .06	0.4 $\pm$ .14	0.0 $\pm$ .00	.6 $\pm$ .16	1.1 $\pm$ .43	0.0 $\pm$ .00	0.2 $\pm$ .11	0.5 $\pm$ .15	0.1 $\pm$ .06	0.9 $\pm$ .17	0.3 $\pm$ .13
<u>Lygodesmia juncea</u>	15	0.0 $\pm$ .00	0.3 $\pm$ .12	0.6 $\pm$ .16	0.2 $\pm$ .09	0.8 $\pm$ .43	0.9 $\pm$ .43	0.1 $\pm$ .06	0.3 $\pm$ .12	0.8 $\pm$ .43	0.1 $\pm$ .08	0.3 $\pm$ .13	0.8 $\pm$ .43
<u>Petalostemum purpureum</u>	12	0.0 $\pm$ .00	0.1 $\pm$ .08	1.8 $\pm$ .79	0.1 $\pm$ .06	1.2 $\pm$ .58	1.1 $\pm$ .58	0.0 $\pm$ .00	0.7 $\pm$ .43	0.9 $\pm$ .58	0.1 $\pm$ .06	0.3 $\pm$ .13	0.5 $\pm$ .42
<u>Juniperus virginiana</u>	11	1.4 $\pm$ .58	0.3 $\pm$ .12	0.1 $\pm$ .08	0.3 $\pm$ .12	0.0 $\pm$ .00	0.0 $\pm$ .00	1.2 $\pm$ .58	0.1 $\pm$ .08	0.1 $\pm$ .08	0.3 $\pm$ .13	0.2 $\pm$ .09	0.1 $\pm$ .06
<u>Echinacea angustifolia</u>	9	0.0 $\pm$ .00	0.2 $\pm$ .11	0.3 $\pm$ .12	0.1 $\pm$ .08	0.1 $\pm$ .06	0.2 $\pm$ .11	0.1 $\pm$ .08	0.2 $\pm$ .09	0.2 $\pm$ .11	0.2 $\pm$ .09	0.3 $\pm$ .13	0.17 $\pm$ .09
<u>Muhlenbergia cuspidata</u>	6	0.0 $\pm$ .00	1.1 $\pm$ 1.06	1.7 $\pm$ 1.72	0.4 $\pm$ .42	0.4 $\pm$ .42	0.4 $\pm$ .42	0.0 $\pm$ .00	0.4 $\pm$ .42	0.5 $\pm$ .42	1.1 $\pm$ 1.06	0.1 $\pm$ .06	0.0 $\pm$ .00
<u>Anemone cylindrica</u>	5	0.2 $\pm$ .09	0.2 $\pm$ .09	0.1 $\pm$ .06	0.2 $\pm$ .11	0.1 $\pm$ .06	0.0 $\pm$ .00	0.6 $\pm$ .42	0.1 $\pm$ .06	0.1 $\pm$ .08	0.1 $\pm$ .06	0.1 $\pm$ .06	0.1 $\pm$ .08
<u>Erigeron strigosus</u>	5	0.1 $\pm$ .06	0.1 $\pm$ .06	0.0 $\pm$ .00	0.1 $\pm$ .06	0.1 $\pm$ .06	0.1 $\pm$ .08	0.1 $\pm$ .06	0.2 $\pm$ .09	0.2 $\pm$ .11	0.0 $\pm$ .00	0.1 $\pm$ .08	0.2 $\pm$ .11

Appendix Table I. Floristics of study plots (continued). Mean percent canopy cover  $\pm$  SE, and relative frequency.

Species:	% Rel. Freq.	Mean percent canopy cover $\pm$ SE											
		1-N	2-N	3-N	1-S	2-S	3-S	1-E	2-E	3-E	1-W	2-W	3-W
<u>Strophostyles</u> <u>leiosperma</u>	4	0.0 + .00	0.1 + .08	0.1 + .06	0.1 + .08	0.2 + .09	0.1 + .08	0.1 + .06	0.0 + .00	0.0 + .00	0.1 + .06	0.2 + .09	0.2 + .11
<u>Dichantheium</u> <u>wilcoxianum</u>	4	0.0 + .00	0.0 + .00	0.1 + .06	0.1 + .08	0.5 + .42	0.0 + .00	0.0 + .00	0.1 + .06	0.0 + .00	0.1 + .06	0.2 + .09	0.9 + .58
<u>Antennaria</u> <u>neglecta</u>	3	.9 + .58	0.0 + .00	.2 + .09	.1 + .06	.1 + .06	.1 + .06	1.1 + 1.06	.5 + .42	.1 + .06	.1 + .06	0.0 + .00	0.0 + .00
<u>Verbena</u> <u>stricta</u>	3	.1 + .06	0.0 + .00	.1 + .06	0.0 + .00	.1 + .06	.1 + .06	.1 + .06	.2 + .11	0.0 + .06	0.0 + .00	.1 + .06	.1 + .06
<u>Calylophus</u> <u>serrulatus</u>	2	0.0 + .00	.1 + .08	.1 + .08	0.0 + .00	.1 + .06	0.0 + .00	0.0 + .00	.1 + .06	.1 + .08	0.0 + .00	0.0 + .00	.2 + .09
<u>Hedeoma</u> <u>hispidum</u>	2	0.0 + .00	0.0 + .00	0.0 + .00	0.0 + .00	.2 + .09	.7 + .56	0.0 + .00	.1 + .08	0.0 + .00	0.0 + .00	.1 + .08	.1 + .06
<u>Euphorbia</u> <u>marginata</u>	2	0.0 + .00	0.0 + .00	.1 + .06	.1 + .08	0.0 + .00	.1 + .06	0.0 + .00	.1 + .06	0.0 + .00	0.0 + .00	.1 + .08	.1 + .06
<u>Cornus</u> <u>drummondii</u>	2	.1 + .06	0.1 + .06	.4 + .42	0.0 + .00	0.0 + .00	0.0 + .00	.1 + .08	.1 + .06	0.0 + .00	0.0 + .00	0.0 + .00	0.0 + .00
<u>Gaura</u> <u>coccinea</u>	1	0.0 + .00	0.0 + .00	.1 + .06	.1 + .06	0.0 + .00	0.0 + .00	0.0 + .00	.1 + .06	.1 + .06	.1 + .06	.1 + .06	0.0 + .00

Appendix Table I. Floristics of study plots (continued). Mean percent canopy cover  $\pm$  SE, and relative frequency.

Species:	% Rel. Freq.	Mean percent canopy cover $\pm$ SE											
		1-N	2-N	3-N	1-S	2-S	3-S	1-E	2-E	3-E	1-W	2-W	3-W
<u>Sisyrinchium campestre</u>	1	0.0 $\pm$ .00	.1 $\pm$ .06	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .08	0.0 $\pm$ .00	.1 $\pm$ .06
<u>Physalis seedling</u>	1	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	.1 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	.1 $\pm$ .06
<u>Opuntia compressa</u>	1	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00
<u>Ulmus spp.</u>	1	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	.1 $\pm$ .06	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Toxicodendron radicans</u>	1	.1 $\pm$ .06	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Amorpha canescens</u>	1	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.4 $\pm$ .42	1.7 $\pm$ 1.72
<u>Rosa arkansana</u>	1	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	.1 $\pm$ .06
<u>Aster oblongifolia</u>	1	0.0 $\pm$ .00	1.1 $\pm$ 1.06	1.1 $\pm$ 1.06	1.1 $\pm$ 1.06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Mirabilis hirsuta</u>	1	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	.1 $\pm$ .06

Appendix Table I. Floristics of study plots (continued). Mean percent canopy cover  $\pm$  SE, and relative frequency.

Species:	% Rel. Freq.	Mean percent canopy cover $\pm$ SE											
		1-N	2-N	3-N	1-S	2-S	3-S	1-E	2-E	3-E	1-W	2-W	3-W
<u>Koeleria pyramidata</u>	tr	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Aster sagittifolia</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Quercus macrocarpa</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.5 $\pm$ .42	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Ribes missouriense</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .08	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Acer negundo</u>	l	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Fraxinus pennsylvanica</u>	tr	.1 $\pm$ .08	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Parthenocissus seedling</u>	tr	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Sorghastrum nutans</u>	tr	0.0 $\pm$ .00	1.7 $\pm$ 1.72	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	1.1 $\pm$ 1.06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Oxalis spp.</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00



Appendix Table I. Floristics of study plots (continued). Mean percent canopy cover  $\pm$  SE, and relative frequency.


Species:	% Rel. Freq.	Mean percent canopy cover $\pm$ SE											
		1-N	2-N	3-N	1-S	2-S	3-S	1-E	2-E	3-E	1-W	2-W	3-W
<u>Kuhnia spp.</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.4 $\pm$ .42	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Ribes spp.</u>	tr	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Symphoricarpos orbiculatus</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06
<u>Apocynum seedling</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Paspalum setaceum</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Oxytropis lambertii</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Achillea millefolium</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00
<u>Festuca octoflora</u>	tr	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	0.0 $\pm$ .00	.1 $\pm$ .06	0.0 $\pm$ .00

Appendix Table II. Significant differences between sites, separated by plot and aspect. Significance is based on Kruskal-Wallis AOV ( $P < .05$ ). 1=full-canopy, 2=canopy-edge, 3=open-canopy.


Species	SUCCESSIONAL PLOTS
<u>Andropogon scoparius</u>	1-North, 1-West
<u>Poa pratensis</u>	2-South, 2-East, 2-West, 3-North 3-West
<u>Bouteloua curtipendula</u>	2-West
<u>Dichanthelium oligosanthos</u> var. <u>scribnerianum</u>	2-South, 3-East
<u>Andropogon gerardii</u>	1-West, 2-West, 3-East

Appendix Table III. Analysis of variance on transformed-percent-cover for Poa pratensis and other combined grasses.

Source of Variation	Sum of Squares	Df	Mean Square	F	P
<u>POA PRATENSIS</u>					
Main Effects	24.226	10	2.423	40.458	0.000
Plot	20.589	2	10.295	171.922	0.000
Aspect	0.596	3	0.199	3.319	0.020
Site	3.012	5	0.602	10.061	0.000
Interactions	2.756	31	0.089	1.485	0.049
Plot and Aspect	0.819	6	0.136	2.279	0.036
Plot and Site	0.610	10	0.061	1.020	0.426
Aspect and Site	1.330	15	0.089	1.481	0.109
Explained	26.982	41	0.658	10.990	0.000
Residual	23.293	389	0.060		
Total	50.275	430	0.117		
<u>OTHER COMBINED GRASSES</u>					
Main Effects	27.562	10	2.756	44.959	0.000
Plot	25.190	2	12.595	205.448	0.000
Aspect	0.795	3	0.265	4.320	0.005
Site	1.553	5	0.311	5.068	0.000
Interactions	3.349	31	0.108	1.762	0.008
Plot and Aspect	1.025	6	0.171	2.787	0.011
Plot and Site	0.950	10	0.095	1.550	0.120
Aspect and Site	1.370	15	0.091	1.490	0.105
Explained	30.911	41	0.754	12.298	0.000
Residual	23.848	389	0.061		
Total	54.759	430	0.127		



Appendix Figure 1. Regression of transformed-percent-cover of combined other grasses in full-canopy plots onto canopy radius.



Appendix Figure 2. Regression of transformed-percent-cover of Poa pratensis in full-canopy plots onto canopy radius.