

Table 2. Averaged scores for each component of the Medical Outcomes Survey Short Form 36 (SF-36) and the Walking Impairment Questionnaire (WIQ) for Peripheral Arterial Disease (PAD) patients and controls. Significance levels are reported for t-tests between PAD and controls.

<i>Questionnaire scale</i>	<i>PAD patients</i>	<i>Patient</i>	<i>Controls</i>	<i>Control</i>	<i>P-values</i>
		<i>n size</i>		<i>n size</i>	
SF-36 Physical Function	41.5 ± 20.1	n=48	95.3 ± 6.2	n=25	<.001
SF-36 Limitations-Physical	33.3 ± 44.0	n=16	95.8 ± 14.4	n=12	<.001
SF-36 Limitations-Emotional	71.1 ± 45.2	n=16	97.2 ± 9.6	n=12	.061
SF-36 Bodily Pain	46.1 ± 27.6	n=48	75.0 ± 12.5	n=25	<.001
SF-36 Energy	51.1 ± 20.0	n=48	85.0 ± 12.0	n=25	.022
SF-36 Social Functioning	75.5 ± 25.0	n=48	93.3 ± 10.0	n=25	<.001
SF-36 Mental Health	75.3 ± 18.8	n=48	94.1 ± 12.5	n=25	.001
SF-36 General Health	53.8 ± 21.1	n=48	81.1 ± 14.2	n=25	<.001
WIQ Pain	52.6 ± 23.6	n=48	99.5 ± 2.5	n=25	<.001
WIQ Distance	55.0 ± 42.3	n=48	99.8 ± 0.6	n=25	<.001
WIQ Speed	48.3 ± 30.4	n=48	96.8 ± 6.6	n=25	<.001
WIQ Stairs	56.5 ± 47.1	n=48	98.1 ± 4.6	n=25	<.001

Data reported as Mean ± SD. Significance level was P < .05.