

Table 3. Correlation matrix between quantitative clinical measures (Ankle brachial index (ABI), absolute claudication distance, initial claudication distance, and self-selected treadmill pace (SS pace)) and the qualitative self-perceived measures of functional status (Medical Outcomes Survey Short Form 36 (SF-36) and the Walking Impairment Questionnaire (WIQ)).

	<i>n size</i>	<i>ABI</i>	<i>Absolute claudication distance</i>	<i>Initial claudication distance</i>	<i>SS pace</i>
SF-36 Physical Function	n=48	.26	.31*	.23	.01
SF-36 Limitations-Physical	n=16	-.01	.38	.15	-.15
SF-36 Limitations-Emotional	n=16	.10	.26	.24	-.05
SF-36 Bodily Pain	n=48	.13	.26	.46*	.07
SF-36 Vitality	n=48	.10	.30*	.26	.06
SF-36 Social Functioning	n=48	.07	.28	.30*	-.10
SF-36 General Mental Health	n=48	.05	.21	.24	.02
SF-36 General Health	n=48	.08	.09	.01	.13
WIQ pain	n=48	.01	.43*	.40*	-.07
WIQ distance	n=48	.29*	.41*	.35*	.01
WIQ speed	n=48	.32*	.39*	.39*	.08
WIQ stairs	n=48	.11	.21	.19	.05

Significant correlations ($P < .05$) are denoted with as asterisk (*).